

DIMENSION-FREE SCALING LAWS FOR SIZE GENERALIZATION: AN EXPONENTIAL SEPARATION

Behrooz Tahmasebi & Melanie Weber

Harvard John A. Paulson School of Engineering and Applied Sciences (SEAS)

Harvard University, Cambridge, MA 02138, USA

{behrooz_tahmasebi, mweber}@seas.harvard.edu

ABSTRACT

Modern machine learning models increasingly operate on variable-size structured data, such as sets and graphs. A central requirement is *size generalization*: the ability of a model trained on inputs of one size to generalize to inputs of a different, potentially much larger, size. Classical statistical learning theory, however, primarily considers fixed input domains, and its guarantees often deteriorate with input dimension. Although extensive work has established when models on sets and graphs can be transferred across input sizes, such results do not characterize the statistical cost of doing so: the sample complexity of size generalization, and its relationship to classical in-domain learning, remain largely unknown. We develop a statistical learning theory of size generalization and establish a sharp exponential separation in sample complexity. Letting n denote the number of training samples, we show that polynomial source decay is insufficient for efficient size generalization: while it yields error $n^{-\Theta(1)}$ and hence polynomial sample complexity for in-domain learning, size-generalization error decays only as $(\log n)^{-\Theta(1)}$, yielding an exponential separation. Thus, polynomial in-domain sample complexity can become exponential across sizes. We then identify an exponential source condition under which size generalization achieves error $n^{-\Theta(1)}$, recovering polynomial sample complexity. These guarantees are independent of input dimension and depend only on the number of training samples, yielding dimension-free scaling laws. We also provide matching minimax lower bounds, establishing optimality. To our knowledge, this work provides the first dimension-free minimax theory of size generalization and a rigorous characterization of when, and at what statistical cost, learning can transfer across input sizes.

1 INTRODUCTION

In many geometric learning problems, the domain is not fixed once and for all. Sets vary in cardinality, graphs vary in the number of nodes, and objects encountered at test time may be larger than those observed during training. The goal is therefore not only to generalize to unseen in-domain samples but also to generalize across domains of different sizes. This desirable property is central to geometric machine learning (Bronstein et al., 2021; Weber, 2025) and is often studied as *size generalization* for sets and graphs (Yehudai et al., 2021; Bevilacqua et al., 2021).

Recent work has made important progress toward understanding learning across sizes, but the available theory remains limited. One line studies transferability through graph limits, graphons, and related stability frameworks (Ruiz et al., 2020; Keriven et al., 2020; Levie et al., 2021; Levin et al., 2025), while another develops learning rules across dimensions using representation stability (Farb, 2014; Levin & Díaz, 2024; Levin & Chandrasekaran, 2023; Díaz et al., 2025). These works offer structural and statistical insights, but leave open a dimension-free minimax characterization of sample complexity across sizes.

The ability to process variable-size inputs does not by itself close this gap. Deep Sets (Zaheer et al., 2017) and graph neural networks (Scarselli et al., 2008), for example, can be applied to sets and graphs of varying sizes and may even satisfy transferability guarantees: under suitable regularity conditions, such as Lipschitz continuity, a fixed model produces similar outputs on objects of different

sizes. But such guarantees do not determine how the transferred error decreases with the number n of training samples, whether this rate is uniform over the training and test sizes, or how it compares with generalization at the training size. We therefore ask: given n samples at size d , can we control the generalization error simultaneously at every target size $D \geq d$, using a number of samples independent of both d and D ?

To address this question, we return to a classical idea. In classical learning theory, it is standard to impose a *source condition* on the desired task, since learning over an arbitrary space is impossible by a standard no-free-lunch argument. The source condition that enables the in-domain generalization error to decrease as $n^{-\Theta(1)}$ is called polynomial source decay. Roughly speaking, it requires the Fourier coefficients of the target function to decay polynomially with their index. This condition is necessary and sufficient for efficient classical in-domain learning. Motivated by this phenomenon, we ask the following question:

Do the conditions for efficient in-domain learning also enable efficient size generalization?

We therefore ask whether polynomial source decay remains sufficient for size generalization. Our main finding is a sharp separation: under polynomial source decay, the size-generalization error decreases only as $(\log n)^{-\Theta(1)}$. Consequently, an accuracy attainable with polynomially many samples for in-domain learning under the same condition may require exponentially many samples when the task must generalize across input sizes.

Table 1: The statistical cost of size generalization.

Source decay	In-domain error	Size-generalization error
Polynomial	$n^{-\Theta(1)}$	$(\log n)^{-\Theta(1)}$
Exponential	$n^{-\Theta(1)}$	$n^{-\Theta(1)}$

Importantly, our resulting guarantees are *dimension-free*: they depend on the number of training samples, but not on the input size d (i.e., either the cardinality of a set or the number of nodes in a graph). In particular, classical transferability errors are typically controlled by the smaller object size, scaling as $d^{-\Theta(1)}$ when $D \geq d$. By contrast, our generalization gap converges to zero with the number of training samples n and is uniform over both d and D , albeit at a slow rate.

To overcome this limitation, we then identify when the exponential cost can be avoided. Under *exponential source decay*, which is stronger than polynomial source decay, we show that the size-generalization error improves from $(\log n)^{-\Theta(1)}$ to $n^{-\Theta(1)}$, recovering polynomial sample complexity. Therefore, our main finding is:

Efficient size generalization requires stronger regularity than in-domain learning; polynomial decay is sufficient for the latter, whereas exponential decay is needed for the former.

Table 1 summarizes this contrast. Exponential decay arises naturally for heat (Gaussian) kernels, analytic functions, and local graph functions.

Finally, we prove matching minimax lower bounds for both sets and graphs. The polynomial-versus-exponential separation is therefore not caused by a particular algorithmic issue. It is an inherent statistical limitation of size generalization over the corresponding source classes. Together, the upper and lower bounds provide a dimension-free minimax characterization of when learning can generalize across input sizes and at what sample cost.

In short, in this paper we make the following contributions:

- We develop dimension-free scaling laws for invariant learning and estimation on sets and graphs, with rates that hold uniformly across input sizes (Section 4 and Theorem 4.1).
- We show that size generalization is fundamentally harder than classical in-domain statistical learning: under polynomial source decay, in-domain error is $n^{-\Theta(1)}$, while size-generalization error is only $(\log n)^{-\Theta(1)}$, implying an exponential separation in sample complexity (Theorem 4.1 and Corollary 4.3).

- We show that under exponential source decay, size-generalization error becomes $n^{-\Theta(1)}$, recovering polynomial sample complexity independently of object size, at the cost of stronger regularity (Theorem 4.1).
- We prove that the rates for both sets and graphs are minimax optimal, establishing that the separation is intrinsic rather than algorithm-dependent (Theorem 4.2).

2 RELATED WORK

Transferability and size generalization. Transferability across graph sizes has been studied through graph limits, graphons, graphops, and higher-order limits (Ruiz et al., 2020; Keriven et al., 2020; Levie et al., 2021; Ruiz et al., 2023; Maskey et al., 2023; Cai & Wang, 2022; Le & Jegelka, 2023; Herbst & Jegelka, 2025). Related work studies learning across set cardinalities and learning-theoretic formulations of graph size generalization (Zweig & Bruna, 2021; Maehara & NT, 2021). These works primarily characterize approximation, consistency, or stability across sizes. In contrast, we ask when an estimator learned at one size achieves statistical convergence rates that are uniform over all test sizes.

Representation stability and any-dimensional learning. Our work builds on representation stability (Farb, 2014), which describes how structured families of representations stabilize across dimensions. Representation stability has recently been used in machine learning to design and analyze architectures for learning under symmetries (Levin & Díaz, 2024), optimization across dimensions (Levin & Chandrasekaran, 2023), graph limits (Levin & Chandrasekaran, 2026), scientific computing and PDEs (Phan et al., 2025), and finite-dimensional kernel learning under invariances (Díaz et al., 2025). The work of Levin et al. (2025), which establishes size-generalization bounds in a broad framework encompassing sets and graphs, is most closely related to ours. Although their framework provides transfer and generalization guarantees across dimensions, the resulting rates depend on dimension-dependent complexity measures, such as covering numbers of the domains. In contrast, we identify conditions under which convergence rates are independent of object size and characterize the tradeoff between invariant feature dimension growth and target approximation. To the best of our knowledge, our work is the first to establish this sharp separation between slow and fast regimes together with matching minimax optimality guarantees. For a broader discussion of related work, see the appendix.

3 PROBLEM STATEMENT

Sets and (attributed) graphs. We study problems indexed by the object size d , the number of elements in a set or nodes in a graph, and fix a feature dimension $r \in \mathbb{N}$. A set is represented by $X = (x_1, \dots, x_d) \in (\mathbb{R}^r)^d \simeq \mathbb{R}^{r \times d}$, while an attributed graph is represented by $X \in \mathbb{R}^{r \times d \times d}$, with r edge and/or node-feature channels. Accordingly, $\mathcal{X}_d$ denotes $\mathbb{R}^{r \times d}$ or $\mathbb{R}^{r \times d \times d}$. These matrix and tensor representations are ordered, but the underlying sets and graphs are not. We equip $\mathcal{X}_d$ with the standard Gaussian probability measure λ_d , under which the vectorized entries are independent $\mathcal{N}(0, 1)$. This measure is permutation invariant and has positive Lebesgue density everywhere, so it can serve as a reference for any distribution with a Lebesgue density; we choose it because Hermite polynomials form a convenient orthogonal basis.

Statistical learning at a fixed size. We consider unsupervised density estimation and supervised regression together. Write $Z_i = X_i$ in the unsupervised setting, where $X_i \stackrel{\text{i.i.d.}}{\sim} \mu_d$ and $d\mu_d = f_d^* d\lambda_d$, and write $Z_i = (X_i, Y_i)$ in the supervised setting, where $(X_i, Y_i) \stackrel{\text{i.i.d.}}{\sim} \rho_d$, $Y_i \in \mathbb{R}$, the X -marginal is μ_d , and $f_d^*(x) = \mathbb{E}[Y \mid X = x]$. Let $\mathcal{H}_d = L^2(\lambda_d)$ for density estimation and $\mathcal{H}_d = L^2(\mu_d)$ for regression, and assume $f_d^* \in \mathcal{F}_d \subseteq \mathcal{H}_d$ is the target function to estimate. For the corresponding loss $\ell_d(Z, f)$, define

$$\hat{L}_{n,d}(f) = \frac{1}{n} \sum_{i=1}^n \ell_d(Z_i, f), \quad L_d(f) = \mathbb{E}[\ell_d(Z, f)].$$

We assume that ℓ_d identifies f_d^* : the target has finite population risk and is the unique minimizer of L_d over $\mathcal{F}_d$ as an element of L^2 . This uniqueness is functional, not parametric. Many parameter

values may represent the same minimizing function, so the assumption is fully compatible with overparameterized models. For example, negative log-likelihood identifies the true density, while squared loss identifies the conditional mean whenever $\mathbb{E}[Y^2] < \infty$; more generally, proper losses identify their associated population targets. Because μ_d or ρ_d is unknown, L_d cannot be evaluated directly, and empirical risk minimization provides its data-driven counterpart.

Although we state the main formulation for scalar invariant targets, the framework also covers permutation-equivariant vector-valued targets after replacing the invariant spaces and feature counts by their equivariant counterparts; Appendix B.2 gives score matching and denoising as examples. This extension is at the level of function spaces: it requires exact norm-controlled transfer, stable equivariant feature growth, and the corresponding source condition, and its rates are governed by the relevant equivariant feature counts.

Permutation invariance. Let S_d be the symmetric group. A permutation $\pi \in S_d$ acts on sets by $(\pi \cdot X)_{:,j} = X_{:,\pi^{-1}(j)}$ and on graphs by $(\pi \cdot X)_{a,i,j} = X_{a,\pi^{-1}(i),\pi^{-1}(j)}$. Because elements and nodes are unlabeled, we assume $\pi \cdot X \stackrel{\text{law}}{=} X$ for density estimation and $(\pi \cdot X, Y) \stackrel{\text{law}}{=} (X, Y)$ for regression. Invariance of μ_d and λ_d , together with uniqueness of the density, gives $f_d^*(\pi \cdot x) = f_d^*(x)$ almost everywhere in the first case; invariance of the conditional law gives the same conclusion for the regression function. Thus $\mathcal{F}_d^{\text{inv}} := \{f \in \mathcal{F}_d : f(\pi \cdot x) = f(x) \text{ a.e. for every } \pi \in S_d\}$, and the invariant empirical risk minimizer is

$$\hat{f}_{n,d}^{\text{inv}} \in \arg \min_{f \in \mathcal{F}_d^{\text{inv}}} \hat{L}_{n,d}(f).$$

Appendix B.1 provides the formal background.

Exact functional transfer across sizes. Let $\mathcal{H}_d^{\text{inv}} \subseteq \mathcal{H}_d$ be the Hilbert space of invariant functions containing $\mathcal{F}_d^{\text{inv}}$, with the corresponding L^2 norm $\|\cdot\|_d$. For $d \leq D$, let

$$T_{d \rightarrow D} : \mathcal{H}_d^{\text{inv}} \longrightarrow \mathcal{H}_D^{\text{inv}}$$

be a linear operator satisfying $T_{d \rightarrow D}(\mathcal{F}_d^{\text{inv}}) \subseteq \mathcal{F}_D^{\text{inv}}$. It maps an entire rule, independently of its parameterization, from size d to size D . We assume coherence, $T_{d \rightarrow d} = \text{Id}$ and $T_{D \rightarrow E} \circ T_{d \rightarrow D} = T_{d \rightarrow E}$ for $d \leq D \leq E$; exact target transfer, $T_{d \rightarrow D} f_d^* = f_D^*$; and norm preservation, $\|T_{d \rightarrow D} f - T_{d \rightarrow D} h\|_D = \|f - h\|_d$. Thus the targets at different sizes are realizations of a single size-consistent rule, with no transfer mismatch or approximation error between sizes. This consistency assumption is standard and essential in the size-generalization literature (Levin et al., 2025).

This is the function-level abstraction of the shared rules used by Deep Sets, GNNs, invariant feature expansions, and graphon representations (Zaheer et al., 2017; Xu et al., 2019; Ruiz et al., 2020; Levin & Díaz, 2024; Levin et al., 2025). The map need not cover every function at size D ; it only carries the same rule to the larger domain while preserving its error scale. Without such a connection, size- d data would contain no information about an arbitrary size- D target. Appendices B.3 and C give formal definitions and concrete constructions.

A concrete picture. An invariant rule can be viewed as a weighted combination of patterns whose meaning does not change with object size. For a set, a pattern might measure an average feature or an interaction among a few elements; for a graph, it might measure a node or edge feature or the frequency of a small subgraph. Let $\psi_{d,j}$ denote the normalized measurement of pattern j on an object of size d . The same index j refers to the same pattern at every size. A rule $\sum_j a_j \psi_{d,j}$ is therefore transferred to size D simply by reusing the learned weights:

$$T_{d \rightarrow D} \left(\sum_j a_j \psi_{d,j} \right) = \sum_j a_j \psi_{D,j}.$$

With a suitable normalization, the squared difference between two rules is exactly the squared difference between their weights. Reusing the weights at a new size therefore preserves prediction error; formally, this is the orthonormal-basis construction and Parseval’s identity. Exact target transfer then has a simple meaning: the true task uses the same pattern weights $(a_j)_j$ at every size. Thus, the assumption does not identify arbitrary functions across dimensions; it only identifies the same invariant rule evaluated on different numbers of elements or nodes. Appendices B.3 and C construct these features explicitly for sets and graphs.

The statistical objective: generalization across size. Training uses only n samples at size d . At any target size $D \geq d$, the learned function is $T_{d \rightarrow D} \hat{f}_{n,d}^{\text{inv}}$. Our goal is a guarantee whose sample complexity is independent of both sizes:

Dimension-free size generalization. For every accuracy $\varepsilon > 0$ and confidence level $1 - \delta$, find $n = n(\varepsilon, \delta)$, independent of d and D , such that, with probability at least $1 - \delta$,

$$\sup_{D \geq d} \|T_{d \rightarrow D} \hat{f}_{n,d}^{\text{inv}} - f_D^*\|_D^2 \leq \varepsilon.$$

The supremum is essential: a single estimator, learned at one size, must remain accurate simultaneously at every larger size. The remainder of the paper characterizes exactly when this is statistically possible and how the required number of samples scales with the regularity of the target.

4 MAIN RESULTS

We now give the dimension-free rates for statistical learning on sets and graphs. The result has two ingredients: representation stability determines how many invariant interactions exist at each order, while target regularity determines how much of the learning task remains beyond that order. Combining them yields a sharp phase transition between nearly parametric and logarithmic size generalization.

4.1 INVARIANT POLYNOMIAL COORDINATES AND REPRESENTATION STABILITY

Recall the domain $\mathcal{X}_d$, target class $\mathcal{F}_d$, and Hilbert space $\mathcal{H}_d$ from Section 3. Thus, $\mathcal{F}_d \subseteq \mathcal{H}_d^{\text{inv}}$, with $\mathcal{H}_d = L^2(\lambda_d)$ for density estimation and $\mathcal{H}_d = L^2(\mu_d)$ for regression.

To organize this infinite-dimensional space, let $\mathcal{V}_{d,k}$ be the span of all invariant polynomials of degree at most k on $\mathcal{X}_d$. Under the standard completeness conditions stated in the appendix, polynomials are dense in $\mathcal{H}_d$; averaging over relabelings then makes invariant polynomials dense in $\mathcal{H}_d^{\text{inv}}$. The polynomial description therefore does not assume that the target itself is a polynomial. It supplies a complete, transfer-compatible hierarchy in which degree records interaction complexity: low degrees describe low-order interactions, while increasing k progressively resolves richer dependencies among elements or nodes.

An orthonormal basis for $\mathcal{V}_{d,k}$ is obtained by applying Gram–Schmidt in $\mathcal{H}_d$, with inner product $\langle \cdot, \cdot \rangle_d$. Orthogonalization changes only the coordinates, not the polynomial span or its dimension. The resulting nested spaces

$$\mathcal{V}_{d,0} \subseteq \mathcal{V}_{d,1} \subseteq \dots \subseteq \mathcal{H}_d^{\text{inv}}$$

are compatible with the exact transfer operators, and representation stability controls their complexity uniformly over object size. Specifically, for $X \in \{\text{set}, \text{graph}\}$, define

$$q_X(k) := \sup_{d \geq 1} \dim(\mathcal{V}_{d,k}).$$

For every fixed k , the dimensions increase to a stable value, so the supremum equals that value and bounds every size, including the pre-stable regime (Farb, 2014).

For sets $X \in \mathbb{R}^{r \times d}$, these are multisymmetric polynomials. Hardy–Ramanujan and Meinardus partition asymptotics, detailed in Appendix B.5, give

$$q_{\text{set}}(k) = \exp(C_r k^\beta + o(k^\beta)), \quad \beta = \frac{r}{r+1}.$$

Here $C_r = \frac{r+1}{r} (r\zeta(r+1))^{1/(r+1)}$, where ζ is the Riemann zeta function. For attributed graphs $X \in \mathbb{R}^{r \times d \times d}$ with at least one edge channel, invariant polynomials are indexed by unlabeled attributed multigraph patterns. The enumeration in Appendix B.7 gives

$$q_{\text{graph}}(k) = \exp(k \log k + O_r(k \log \log k)).$$

For fixed r , the graph feature dimension depends on r only through the lower-order term $O_r(k \log \log k)$. Consequently, r does not enter the first-order graph asymptotics or the graph rates below. Both feature counts are uniform in the number of elements or nodes, but the number of interactions grows rapidly with degree and substantially faster for graphs.

Table 2: Minimax-optimal in-domain and dimension-free size-generalization rates.

Source decay	In-domain	Sets	Graphs
Polynomial, $k^{-\tau}$	Sets: $n^{-\tau/(\tau+rd)}$ Graphs: $n^{-\tau/(\tau+rd^2)}$	$(\log n)^{-\tau(r+1)/r}$	$(\log n / \log \log n)^{-\tau}$
Exponential, e^{-ak^ν}	Both: $n^{-1+o(1)}$	$n^{-1+o(1)}$ if $\nu > \frac{r}{r+1}$	$n^{-1+o(1)}$ if $\nu > 1$

4.2 POLYNOMIAL AND EXPONENTIAL SOURCE DECAY

Let $\mathcal{F}_{d,k} = \mathcal{V}_{d,k}$ for regression and $\mathcal{F}_{d,k} = \mathcal{F}_d^{\text{inv}} \cap \mathcal{V}_{d,k}$ for density estimation, so density approximants remain nonnegative and normalized. Define the best order- k approximation error and its uniform envelope by

$$B_d(k) := \inf_{f \in \mathcal{F}_{d,k}} \|f_d^* - f\|_d^2, \quad B(k) := \sup_{d \geq 1} B_d(k).$$

For regression, the optimizer is the orthogonal projection $\Pi_{d,k} f_d^*$; hence $B(k)$ is the squared energy in coordinates of degree greater than k , uniformly over d . The density definition is its admissible analogue. In both cases, this is an invariant Fourier or spectral source condition: the basis is complete, while coefficient decay quantifies how strongly the target depends on high-order interactions.

We use the two source conditions introduced in the paper:

$$\begin{aligned} \text{polynomial source decay: } B(k) &\lesssim k^{-\tau}, & \tau > 0, \\ \text{exponential source decay: } B(k) &\lesssim \exp(-ak^\nu), & a, \nu > 0. \end{aligned}$$

Polynomial decay is the analogue of finite smoothness: increasingly complex interactions contribute progressively less, but remain present at every order. Exponential decay describes substantially stronger regularity. The exponent ν determines whether this decay is fast enough to overcome the growth of invariant interactions.

4.3 DIMENSION-FREE SCALING LAWS

For an interaction order k , let $f_{d,k}^\circ$ minimize the population risk over $\mathcal{F}_{d,k}$, and let $\hat{f}_{n,d,k}^{\text{inv}}$ be its empirical counterpart. We impose standard, primitive regularity conditions on this finite-dimensional problem. In orthonormal coordinates on $\mathcal{V}_{d,k}$, the population approximation satisfies $\|f_{d,k}^\circ - f_d^*\|_d^2 \lesssim B_d(k)$; the empirical loss has curvature at least $\alpha > 0$ on the relevant neighborhood; and the centered empirical score at $f_{d,k}^\circ$ is sub-Gaussian with parameter at most σ^2 . The constants α and σ are uniform in d and k . Here, curvature means that moving the fitted function by h increases the loss by at least $\alpha \|h\|_d^2/2$, up to its linear term; it rules out statistically invisible directions.

These conditions are directly checkable for the canonical losses in our framework. For squared-loss regression,

$$L_d(f) - L_d(f_d^*) = \|f - f_d^*\|_{L^2(\mu_d)}^2,$$

so approximation is exactly measured by $B_d(k)$. Curvature follows from concentration of the empirical Gram matrix, and score concentration follows from sub-Gaussian regression noise and whitened invariant features. For density estimation by negative log-likelihood, the Hessian is uniformly positive and the likelihood is locally equivalent to squared $L^2(\lambda_d)$ error when candidate densities are bounded above and away from zero; bounded or sub-Gaussian invariant scores then give the same concentration. Regular proper losses satisfy the analogous calibration and Hessian conditions. Appendix F derives the finite-dimensional estimation bound from these assumptions rather than postulating it.

Table 2 compares our dimension-free rates with classical in-domain learning. At a fixed size d , the number of degree-at-most- k invariant polynomial features grows as k^{rd} for sets and k^{rd^2} for graphs under our tensor representation; invariance under the finite group S_d changes constants but not these exponents. Constants and lower-order factors are suppressed throughout.

At and below the exponential thresholds, the rates interpolate between the two rows of the table. The exact critical and subcritical expressions are derived in Appendix D; the main distinction is whether exponential source decay is fast enough to recover a nearly parametric rate.

Theorem 4.1 (Dimension-free scaling laws for sets and graphs). *Assume exact isometric transfer, representation stability, compatibility $T_{d \rightarrow D}(\mathcal{F}_{d,k}) \subseteq \mathcal{F}_{D,k}$, and the uniform loss regularity above. For any fixed confidence level, there are invariant empirical estimators, trained using only size- d data, whose squared error holds simultaneously for every $D \geq d$ and satisfies the following rates, independently of d and D :*

(i) *If $B(k) \lesssim k^{-\tau}$, the rate is*

$$(\log n)^{-\tau(r+1)/r} \quad \text{for sets,} \quad \left(\frac{\log n}{\log \log n} \right)^{-\tau} \quad \text{for graphs.}$$

(ii) *If $B(k) \lesssim e^{-ak^\nu}$, the set rate, with $\beta = r/(r+1)$, is*

$$\begin{cases} n^{-1+o(1)}, & \nu > \beta, \\ n^{-a/(a+C_r)+o(1)}, & \nu = \beta, \\ \exp\left[-aC_r^{-\nu/\beta}(\log n)^{\nu/\beta}(1+o(1))\right], & 0 < \nu < \beta, \end{cases}$$

while the graph rate is $n^{-1+o(1)}$ for $\nu > 1$ and

$$\exp\left[-a \left(\frac{\log n}{\log \log n} \right)^\nu (1+o(1))\right] \quad \text{for } 0 < \nu \leq 1.$$

The statements apply to squared-loss regression, regular likelihood density estimation, and other proper losses satisfying the stated calibration, curvature, and score conditions.

The theorem is uniform in a strong sense: one event controls the transferred estimator at every larger deployment size, without a union bound over D . Representation stability removes object size from the rate, but it does not make all invariant interactions equally easy to learn. For polynomial source classes, nonnegligible structure persists across the rapidly expanding catalogue of stable interactions, forcing logarithmic convergence. Exponential decay suppresses those high-order interactions fast enough to recover a power law, with different thresholds for sets and graphs because their invariant coordinates grow at different rates.

4.4 WHY CLASSICAL IN-DOMAIN POWER LAWS CHANGE

In classical fixed-domain nonparametric learning, the number of features up to resolution k often grows polynomially, say $q(k) \asymp k^s$. Polynomial source decay $B(k) \asymp k^{-\tau}$ then produces the familiar power law $n^{-\tau/(\tau+s)}$. Size generalization on sets and graphs is different. Uniformity over every larger size exposes all invariant interactions, whose stable counts grow as

$$\exp(C_r k^{r/(r+1)}) \quad \text{and} \quad \exp(k \log k),$$

respectively. Consequently, the same polynomial source decay yields only logarithmic convergence in n . A power law is recovered only when the interaction tail has sufficiently fast exponential source decay. This is a statistical effect: it remains even when transfer is exact and optimization is ideal.

To determine whether these rates reflect a particular estimator or an inherent limit of the problem, let $\mathfrak{F}_X(B)$ contain all size-consistent targets of type $X \in \{\text{set}, \text{graph}\}$ with source envelope B . We compare against every measurable procedure, including procedures that do not use polynomial truncation and are not themselves invariant. Its worst-case risk is

$$\mathfrak{R}_n(X, B) := \inf_{\hat{f}} \sup_{d \geq 1} \sup_{f^* \in \mathfrak{F}_X(B)} \mathbb{E} \left[\sup_{D \geq d} \|\hat{f}_D - f_D^*\|_D^2 \right].$$

For the lower bound, the source class must genuinely contain the invariant directions counted by representation stability. Concretely, at every order k , we require a stable-size regression or density submodel with $q_X(k)$ orthogonal local perturbations whose neighboring data distributions have quadratic KL divergence. Gaussian-noise regression and bounded local density or exponential-family perturbations satisfy this condition; Appendix E states and verifies the precise construction.

Theorem 4.2 (Minimax optimality). *Under the preceding local richness condition, every rate in Theorem 4.1 is minimax optimal up to constants and the displayed lower-order factors. In particular, for polynomial source decay,*

$$\mathfrak{R}_n(\text{set}, k^{-\tau}) \asymp (\log n)^{-\tau(r+1)/r}, \quad \mathfrak{R}_n(\text{graph}, k^{-\tau}) \asymp \left(\frac{\log n}{\log \log n} \right)^{-\tau}.$$

For exponential source decay, the minimax risks have the same supercritical, critical, and subcritical regimes stated in Theorem 4.1. These lower bounds hold already at a suitable stable training size and therefore continue to hold when the loss takes a supremum over all $D \geq d$.

The theorem rules out improvements by a different architecture, regularizer, optimization method, or transfer rule: the infimum in $\mathfrak{R}_n$ ranges over all estimators. It also explains the role of exact transfer. Exact transfer removes cross-size mismatch, but it cannot create information about the many statistically distinguishable invariant directions that remain uncertain from n observations. The logarithmic regime is therefore not an artifact of polynomial regression or of the proof technique; it is the optimal statistical behavior of the entire source class.

Corollary 4.3 (Exponential separation in sample complexity). *Fix the confidence level and suppose $B(k) \asymp k^{-\tau}$. Up to constants and lower-order factors, achieving squared error at most ε at a fixed size requires*

$$n_{\text{in}}(\varepsilon) = \begin{cases} \varepsilon^{-(\tau+rd)/\tau}, & \text{sets,} \\ \varepsilon^{-(\tau+rd^2)/\tau}, & \text{graphs,} \end{cases}$$

whereas uniform size generalization requires

$$n_{\text{size}}(\varepsilon) = \begin{cases} \exp\left(\Theta\left(\varepsilon^{-\frac{r}{\tau(r+1)}}\right)\right), & \text{sets,} \\ \exp\left(\Theta\left(\varepsilon^{-1/\tau} \log(1/\varepsilon)\right)\right), & \text{graphs.} \end{cases}$$

Thus the same polynomial source class has polynomial sample complexity in-domain but exponential sample complexity when one estimator must transfer uniformly to all larger sizes.

4.5 DISCUSSION

Our results are stated at the level of function classes rather than a particular neural parameterization. Consequently, they are compatible with permutation-equivariant graph architectures, including graph Transformers, when a permutation-invariant readout produces a graph-level prediction. Such an architecture can parameterize an estimator in our graph setting, but its symmetry alone does not imply the rates above. The cross-size target compatibility, exact norm-controlled transfer, stable feature growth, and source-decay conditions must still be verified. We therefore do not claim that every graph Transformer size-generalizes at these rates; rather, the theorems characterize the statistical behavior of those invariant graph-learning problems that satisfy the stated conditions.

This distinction separates three properties that are often conflated. First, an architecture may be *applicable* to objects of different sizes. Second, a fixed predictor may be *transferable*, meaning that its outputs remain stable as size changes. Third, a predictor learned from finite data may achieve *statistical size generalization*, meaning that its error at unseen sizes decreases at a controlled rate as the number of training samples grows. The first two properties make cross-size prediction possible, but neither determines its sample complexity. Our results address the third property.

The minimax bounds also clarify what would be required to improve the rates. In the polynomial source class, no alternative architecture, regularizer, optimizer, or transfer rule can uniformly replace the logarithmic rate by a power law. Faster rates are possible only by imposing additional structure that reduces the effective class, such as faster source decay or a smaller family of transferable interactions. Thus the exponential separation is not a statement that all individual tasks are hard; it states that polynomial source decay alone is too weak to guarantee efficient size generalization uniformly over the class.

Finally, exact transfer isolates the statistical difficulty studied here by removing cross-size approximation mismatch. With approximate transfer, an additional error term would remain even with unlimited data. Likewise, problems with different symmetries require their own stable feature counts and need

not exhibit the same thresholds. These are natural directions for extending the present theory, but they are distinct from the finite-sample phenomenon characterized by our results.

Altogether, the main message of this paper is:

Exact transfer enables size generalization, but target regularity determines its statistical efficiency. Fast exponential source decay yields nearly parametric rates, whereas polynomial decay yields minimax-optimal logarithmic rates.

5 CONCLUSION

We developed a statistical theory of size generalization for supervised regression and unsupervised density estimation on sets and graphs. By combining exact transfer, representation stability, and source decay in invariant polynomial coordinates, we obtained rates that are uniform over both the training and deployment sizes. Polynomial source decay leads to minimax-optimal logarithmic rates, producing an exponential separation from classical in-domain power laws. Sufficiently fast exponential source decay overcomes invariant feature growth and recovers nearly parametric size generalization, with sharp thresholds that differ between sets and graphs. These results show that architectural transferability alone does not determine the statistical efficiency of size generalization. The decisive interaction is between the growth of transferable invariant representations and the decay of high-order structure in the target. The matching minimax bounds establish that this phase transition is intrinsic rather than an artifact of the estimator or analysis.

AI USE STATEMENT

Large language model tools (ChatGPT 5.6 Sol and Claude Fable 5.1) were used primarily for copyediting, including improvements to grammar, wording, clarity, and $\LaTeX$ presentation. The research direction, core ideas, theoretical results, and experimental design are original contributions of the authors. The authors carefully reviewed and verified all technical content and take full responsibility for the final manuscript.

REFERENCES

- Matan Atzmon, Koki Nagano, Sanja Fidler, Sameh Khamis, and Yaron Lipman. Frame averaging for equivariant shape space learning. In *IEEE Conference on Computer Vision and Pattern Recognition (CVPR)*, 2022. 15
- Beatrice Bevilacqua, Yangze Zhou, and Bruno Ribeiro. Size-invariant graph representations for graph classification extrapolations. In *Int. Conference on Machine Learning (ICML)*, 2021. 1
- Benjamin Bloem-Reddy and Yee Whye Teh. Probabilistic symmetries and invariant neural networks. *Journal of Machine Learning Research*, 21(90):1–61, 2020. 15
- Alexander Bogatskiy, Brandon Anderson, Jan Offermann, Marwah Roussi, David Miller, and Risi Kondor. Lorentz group equivariant neural network for particle physics. In *Int. Conference on Machine Learning (ICML)*, 2020. 15
- Pietro Bongini, Monica Bianchini, and Franco Scarselli. Molecular generative graph neural networks for drug discovery. *Neurocomputing*, 450:242–252, 2021. 15
- Michael M Bronstein, Joan Bruna, Taco Cohen, and Petar Veličković. Geometric deep learning: Grids, groups, graphs, geodesics, and gauges. *arXiv preprint arXiv:2104.13478*, 2021. 1
- Davide Buffelli, Pietro Liò, and Fabio Vandin. Sizeshiftreg: a regularization method for improving size-generalization in graph neural networks. In *Advances in Neural Information Processing Systems (NeurIPS)*, 2022. 15
- Chen Cai and Yusu Wang. Convergence of invariant graph networks. In *Int. Conference on Machine Learning (ICML)*, pp. 2457–2484, 2022. 3

- Andrea Caponnetto and Ernesto De Vito. Optimal rates for the regularized least-squares algorithm. *Foundations of Computational mathematics*, 7(3):331–368, 2007. 15
- Ziyu Chen, Markos Katsoulakis, Luc Rey-Bellet, and Wei Zhu. Sample complexity of probability divergences under group symmetry. In *Int. Conference on Machine Learning (ICML)*, 2023. 15
- Xu Chu, Yujie Jin, Xin Wang, Shanghang Zhang, Yasha Wang, Wenwu Zhu, and Hong Mei. Wasserstein barycenter matching for graph size generalization of message passing neural networks. In *Int. Conference on Machine Learning (ICML)*, 2023. 15
- Ching-Yao Chuang and Stefanie Jegelka. Tree mover’s distance: Bridging graph metrics and stability of graph neural networks. In *Advances in Neural Information Processing Systems (NeurIPS)*, 2022. 15
- Persi Diaconis and Svante Janson. Graph limits and exchangeable random graphs. *Rendiconti di Matematica, Serie VII*, 28:33–61, 2008. 18
- Mateo Díaz, Dmitriy Drusvyatskiy, Jack Kendrick, and Rekha R Thomas. Invariant kernels: Rank stabilization and generalization across dimensions. *arXiv preprint arXiv:2502.01886*, 2025. 1, 3
- Nadav Dym, Hannah Lawrence, and Jonathan W Siegel. Equivariant frames and the impossibility of continuous canonicalization, 2024. 15
- Bryn Elesedy. Provably strict generalisation benefit for invariance in kernel methods. In *Advances in Neural Information Processing Systems (NeurIPS)*, 2021. 15
- Benson Farb. Representation stability. *arXiv preprint arXiv:1404.4065*, 2014. 1, 3, 5
- Vikas Garg, Stefanie Jegelka, and Tommi Jaakkola. Generalization and representational limits of graph neural networks. In *Int. Conference on Machine Learning (ICML)*, 2020. 15
- Daniel Herbst and Stefanie Jegelka. Higher-order graphon neural networks: Approximation and cut distance. In *Int. Conference on Learning Representations (ICLR)*, 2025. 3
- Zheng Huang, Qihui Yang, Dawei Zhou, and Yujun Yan. Enhancing size generalization in graph neural networks through disentangled representation learning. In *Int. Conference on Machine Learning (ICML)*, 2024. 15
- Haotian Ju, Dongyue Li, Aneesh Sharma, and Hongyang R Zhang. Generalization in graph neural networks: Improved pac-bayesian bounds on graph diffusion. In *Int. Conference on Artificial Intelligence and Statistics (AISTATS)*, 2023. 15
- Sékou-Oumar Kaba, Arnab Kumar Mondal, Yan Zhang, Yoshua Bengio, and Siamak Ravanbakhsh. Equivariance with learned canonicalization functions. In *Int. Conference on Machine Learning (ICML)*, 2023. 15
- Nicolas Keriven, Alberto Bietti, and Samuel Vaiter. Convergence and stability of graph convolutional networks on large random graphs. In *Advances in Neural Information Processing Systems (NeurIPS)*, volume 33, pp. 21512–21523, 2020. 1, 3
- Thien Le and Stefanie Jegelka. Limits, approximation and size transferability for gnns on sparse graphs via graphops. In *Advances in Neural Information Processing Systems (NeurIPS)*, 2023. 3
- Ron Levie. A graphon-signal analysis of graph neural networks. *Advances in Neural Information Processing Systems*, 36:64482–64525, 2023. 15
- Ron Levie, Wei Huang, Lorenzo Bucci, Michael Bronstein, and Gitta Kutyniok. Transferability of spectral graph convolutional neural networks. *Journal of Machine Learning Research*, 22(272): 1–59, 2021. 1, 3
- Eitan Levin and Venkat Chandrasekaran. Free descriptions of convex sets. *arXiv preprint arXiv:2307.04230*, 2023. 1, 3
- Eitan Levin and Venkat Chandrasekaran. Limits of weighted graphs via random quotients. *arXiv:2512.23149*, 2026. 3

- Eitan Levin and Mateo Díaz. Any-dimensional equivariant neural networks. In *Int. Conference on Artificial Intelligence and Statistics (AISTATS)*, 2024. 1, 3, 4, 17
- Eitan Levin, Yuxin Ma, Mateo Díaz, and Soledad Villar. On transferring transferability: Towards a theory for size generalization. In *Advances in Neural Information Processing Systems (NeurIPS)*, 2025. 1, 3, 4
- Xiao Li, Li Sun, Mengjie Ling, and Yan Peng. A survey of graph neural network based recommendation in social networks. *Neurocomputing*, 549:126441, 2023. 15
- Renjie Liao, Raquel Urtasun, and Richard Zemel. A pac-bayesian approach to generalization bounds for graph neural networks. In *Int. Conference on Learning Representations (ICLR)*, 2020. 15
- Yuchao Lin, Jacob Helwig, Shurui Gui, and Shuiwang Ji. Equivariance via minimal frame averaging for more symmetries and efficiency. In *Int. Conference on Machine Learning (ICML)*, 2024. 15
- George Ma, Yifei Wang, Derek Lim, Stefanie Jegelka, and Yisen Wang. A canonicalization perspective on invariant and equivariant learning. In *Advances in Neural Information Processing Systems (NeurIPS)*, 2024. 15
- Takanori Maehara and Hoang NT. Learning on random balls is sufficient for estimating (some) graph parameters. In *Advances in Neural Information Processing Systems (NeurIPS)*, 2021. 3
- Haggai Maron, Heli Ben-Hamu, Nadav Shamir, and Yaron Lipman. Invariant and equivariant graph networks. In *Int. Conference on Learning Representations (ICLR)*, 2019. 15
- Sohir Maskey, Ron Levie, Yunseok Lee, and Gitta Kutyniok. Generalization analysis of message passing neural networks on large random graphs. *Advances in neural information processing systems*, 35:4805–4817, 2022. 15
- Sohir Maskey, Ron Levie, and Gitta Kutyniok. Transferability of graph neural networks: an extended graphon approach. *Applied and Computational Harmonic Analysis*, 63:48–83, 2023. 3
- Sohir Maskey, Gitta Kutyniok, and Ron Levie. Generalization bounds for message passing networks on mixture of graphons. *SIAM Journal on Mathematics of Data Science*, 7(2):802–825, 2025. 15
- Christopher Morris, Floris Geerts, Jan Tönshoff, and Martin Grohe. Wl meet vc. In *Int. Conference on Machine Learning (ICML)*, 2023. 15
- Quynh T Nguyen, Louis Schatzki, Paolo Braccia, Michael Ragone, Patrick J Coles, Frederic Sauvage, Martin Larocca, and Marco Cerezo. Theory for equivariant quantum neural networks. *PRX Quantum*, 5(2):020328, 2024. 15
- Mircea Petrache and Shubhendu Trivedi. Approximation-generalization trade-offs under (approximate) group equivariance. In *Advances in Neural Information Processing Systems (NeurIPS)*, 2023. 15
- Trung V Phan, George A Kevrekidis, Soledad Villar, Yannis G Kevrekidis, and Juan M Bello-Rivas. Towards coordinate-and dimension-agnostic machine learning for partial differential equations. *arXiv preprint arXiv:2505.16549*, 2025. 3
- Omri Puny, Matan Atzmon, Edward J Smith, Ishan Misra, Aditya Grover, Heli Ben-Hamu, and Yaron Lipman. Frame averaging for invariant and equivariant network design. In *Int. Conference on Learning Representations (ICLR)*, 2022. 15
- Levi Rauchwerger, Stefanie Jegelka, and Ron Levie. Generalization, expressivity, and universality of graph neural networks on attributed graphs. In *Int. Conference on Learning Representations (ICLR)*, 2025. 15
- Luana Ruiz, Luiz Chamon, and Alejandro Ribeiro. Graphon neural networks and the transferability of graph neural networks. In *Advances in Neural Information Processing Systems (NeurIPS)*, volume 33, pp. 1702–1712, 2020. 1, 3, 4, 18
- Luana Ruiz, Luiz FO Chamon, and Alejandro Ribeiro. Transferability properties of graph neural networks. *IEEE Transactions on Signal Processing*, 71:3474–3489, 2023. 3

- Victor Garcia Satorras, Emiel Hoogetboom, and Max Welling. E(n) equivariant graph neural networks. In *Int. Conference on Machine Learning (ICML)*, 2021. 15
- Franco Scarselli, Marco Gori, Ah Chung Tsoi, Markus Hagenbuchner, and Gabriele Monfardini. The graph neural network model. *IEEE transactions on neural networks*, 20(1):61–80, 2008. 1
- Franco Scarselli, Ah Chung Tsoi, and Markus Hagenbuchner. The Vapnik–Chervonenkis dimension of graph and recursive neural networks. *Neural Networks*, 108:248–259, 2018. 15
- Zakhar Shumaylov, Peter Zaika, James Rowbottom, Ferdia Sherry, Melanie Weber, and Carola-Bibiane Schönlieb. Lie algebra canonicalization: Equivariant neural operators under arbitrary Lie groups. In *Int. Conference on Learning Representations (ICLR)*, 2025. 15
- Ashkan Soleymani, Behrooz Tahmasebi, Stefanie Jegelka, and Patrick Jaillet. A robust kernel statistical test of invariance: Detecting subtle asymmetries. In *Int. Conference on Artificial Intelligence and Statistics (AISTATS)*, 2025a. 15
- Ashkan Soleymani, Behrooz Tahmasebi, Stefanie Jegelka, and Patrick Jaillet. Learning with exact invariances in polynomial time. In *Int. Conference on Machine Learning (ICML)*, 2025b. 15
- Ashkan Soleymani, Behrooz Tahmasebi, Patrick Jaillet, and Stefanie Jegelka. Efficient learning and symmetry discovery under exact invariances. In *Conference on Learning Theory (COLT)*, 2026a. 15
- Ashkan Soleymani, Behrooz Tahmasebi, Patrick Jaillet, and Stefanie Jegelka. A unified framework for statistical testing of invariance. In *The ICML 2026 Workshop on Hypothesis Testing*, 2026b. 15
- Behrooz Tahmasebi and Stefanie Jegelka. The exact sample complexity gain from invariances for kernel regression. In *Advances in Neural Information Processing Systems (NeurIPS)*, 2023. 15
- Behrooz Tahmasebi and Stefanie Jegelka. Sample complexity bounds for estimating probability divergences under invariances. In *Int. Conference on Machine Learning (ICML)*, 2024. 15
- Behrooz Tahmasebi and Stefanie Jegelka. Regularity in canonicalized models: A theoretical perspective. In *Int. Conference on Artificial Intelligence and Statistics (AISTATS)*, 2025a. 15
- Behrooz Tahmasebi and Stefanie Jegelka. Generalization bounds for canonicalization: A comparative study with group averaging. In *Int. Conference on Learning Representations (ICLR)*, 2025b. 15
- Behrooz Tahmasebi and Melanie Weber. Achieving approximate symmetry is exponentially easier than exact symmetry. In *Int. Conference on Learning Representations (ICLR)*, 2026a. 15
- Behrooz Tahmasebi and Melanie Weber. Adaptive symmetry discovery for dynamical system identification. In *Int. Conference on Machine Learning (ICML)*, 2026b. 15
- Behrooz Tahmasebi and Melanie Weber. Sparse data augmentation for optimization with provable guarantees. In *IEEE Conference on Decision and Control (CDC)*, 2026c. 15
- Behrooz Tahmasebi, Derek Lim, and Stefanie Jegelka. The power of recursion in graph neural networks for counting substructures. In *Int. Conference on Artificial Intelligence and Statistics (AISTATS)*, 2023. 15
- Behrooz Tahmasebi, Melanie Weber, and Stefanie Jegelka. Data augmentation: A fourier analysis perspective. In *Conference on Learning Theory (COLT)*, 2026. 15
- Antonis Vasileiou, Ben Finkelshtein, Floris Geerts, Ron Levie, and Christopher Morris. Covered forest: Fine-grained generalization analysis of graph neural networks. In *Int. Conference on Machine Learning (ICML)*, 2025a. 15
- Antonis Vasileiou, Stefanie Jegelka, Ron Levie, and Christopher Morris. Survey on generalization theory for graph neural networks. *arXiv preprint arXiv:2503.15650*, 2025b. 15
- Petar Veličković, Christos Perivolaropoulos, Federico Barbero, and Razvan Pascanu. Softmax is not enough (for sharp size generalisation). In *Int. Conference on Machine Learning (ICML)*, 2025. 15

- Edward Wagstaff, Fabian Fuchs, Martin Engelcke, Ingmar Posner, and Michael A Osborne. On the limitations of representing functions on sets. In *Int. Conference on Machine Learning (ICML)*, 2019. 15
- Dian Wang, Robin Walters, and Robert Platt. $SO(2)$ -equivariant reinforcement learning. In *Int. Conference on Learning Representations (ICLR)*, 2022. 15
- Rui Wang, Robin Walters, and Rose Yu. Incorporating symmetry into deep dynamics models for improved generalization. In *Int. Conference on Learning Representations (ICLR)*, 2021. 15
- Zhiyang Wang, Juan Cervino, and Alejandro Ribeiro. A manifold perspective on the statistical generalization of graph neural networks. In *Int. Conference on Machine Learning (ICML)*, 2025. 15
- Melanie Weber. Geometric machine learning. *Wiley Online Library*, 2025. 1
- Keyulu Xu, Weihua Hu, Jure Leskovec, and Stefanie Jegelka. How powerful are graph neural networks? In *Int. Conference on Learning Representations (ICLR)*, 2019. 4, 15, 17
- Gilad Yehudai, Ethan Fetaya, Eli Meir, Gal Chechik, and Haggai Maron. From local structures to size generalization in graph neural networks. In *Int. Conference on Machine Learning (ICML)*, 2021. 1
- Manzil Zaheer, Satwik Kottur, Siamak Ravanbakhsh, Barnabas Poczos, Russ R Salakhutdinov, and Alexander J Smola. Deep sets. In *Advances in Neural Information Processing Systems (NeurIPS)*, 2017. 1, 4, 15, 17
- Haobo Zhang, Yicheng Li, Weihao Lu, and Qian Lin. Optimal rates of kernel ridge regression under source condition in large dimensions. *Journal of Machine Learning Research*, 26(219):1–63, 2025. 15
- Aaron Zweig and Joan Bruna. A functional perspective on learning symmetric functions with neural networks. In *Int. Conference on Machine Learning (ICML)*, 2021. 3

APPENDIX CONTENTS

A Additional Related Work	15
B Function-Level Symmetry, Statistical Learning, and Transfer	15
B.1 Group actions on data and functions	16
B.2 Examples of statistical learning objectives	16
B.3 Exact functional transfer across sizes	17
B.4 Invariant polynomial function spaces	18
B.5 Set-valued data and multisymmetric polynomials	18
B.6 Hermite features for set-valued data	19
B.7 Attributed graphs	20
C Concrete Exact Transfers for Sets and Graphs	21
C.1 Sets	21
C.2 Attributed graphs	22
C.3 What the exact-transfer assumption asserts	23
D Proof of Theorem 4.1	23
D.1 Finite-dimensional estimation and transfer	23
D.2 Set rates	24
D.3 Graph rates	24
E Proof of Theorem 4.2	25
E.1 Minimax risk and local invariant submodels	25
E.2 Assouad lower bound	25
E.3 Evaluation of the lower bound	26
F Auxiliary finite-dimensional estimation bound	26
F.1 Restricted gradients on the invariant model	26
F.2 Proof of the estimation bound	27
F.3 Proof of the transfer bound	28
G Equivariant Field Specialization for Sets	28
H Equivariant Field Specialization for Graphs	29
I Experiments	29
I.1 Subgraph counting	29
I.2 Equivariant field transfer on weighted graphs	30
J Notation	32

A ADDITIONAL RELATED WORK

Learning under invariances. Symmetry and invariance are central principles in geometric machine learning. Deep Sets provide a classical architecture for sets (Zaheer et al., 2017), while graph neural networks and higher-order invariant and equivariant architectures are widely used for graphs (Xu et al., 2019; Maron et al., 2019; Satorras et al., 2021). Probabilistic symmetry provides a distributional foundation for invariant and equivariant models (Bloem-Reddy & Teh, 2020), while complementary work studies limitations of set representations (Wagstaff et al., 2019) and the ability of graph networks to count substructures (Tahmasebi et al., 2023). Beyond architectures that hard-code invariance, architecture-agnostic approaches include canonicalization (Ma et al., 2024; Kaba et al., 2023; Dym et al., 2024; Shumaylov et al., 2025; Tahmasebi & Jegelka, 2025a) and frame averaging (Puny et al., 2022; Atzmon et al., 2022; Lin et al., 2024). Although symmetry must be discovered in some settings, either adaptively (Tahmasebi & Weber, 2026b) or via statistical tests (Soleymani et al., 2025a; 2026b), here we focus on the case of known symmetries. Applications of learning under invariances span reinforcement learning (Wang et al., 2022), physics (Bogatskiy et al., 2020; Nguyen et al., 2024), drug discovery (Bongini et al., 2021), and social networks (Li et al., 2023). Our work is complementary to these modeling approaches: rather than proposing a new architecture, we study when invariant statistical estimation can transfer across dimensions.

Methods and limitations for size generalization. Beyond the transferability theory discussed in the main text, several works develop model-specific approaches to graph size generalization through Wasserstein barycenter matching, size-shift regularization, or disentangled representations (Chu et al., 2023; Buffelli et al., 2022; Huang et al., 2024). Graph metrics have also been used to relate structural shifts to GNN stability (Chuang & Jegelka, 2022). Other results identify architectural limitations to sharp size generalization (Veličković et al., 2025) or derive graph-size-dependent statistical guarantees from a manifold perspective (Wang et al., 2025). These works clarify mechanisms that help or hinder generalization across sizes. Our focus is instead on size-uniform sample complexity and matching minimax rates for invariant estimation.

Generalization and sample complexity under symmetry. The statistical benefits of symmetry have been widely studied in learning theory. Prior work has shown that invariance can reduce sample complexity in supervised learning and related estimation problems (Tahmasebi & Jegelka, 2023; Elesedy, 2021; Wang et al., 2021; Petrache & Trivedi, 2023; Tahmasebi & Jegelka, 2025b), with extensions to estimating probability divergences (Chen et al., 2023; Tahmasebi & Jegelka, 2024). More broadly, the study of symmetry and scaling laws extends to comparisons between approximate and exact symmetry (Tahmasebi & Weber, 2026a), data augmentation via Fourier analysis on groups (Tahmasebi et al., 2026; Tahmasebi & Weber, 2026c), and statistical-computational trade-offs (Soleymani et al., 2025b; 2026a). Generalization bounds for graph neural networks have also been developed using Rademacher complexity (Garg et al., 2020), VC dimension (Scarselli et al., 2018; Morris et al., 2023), and PAC-Bayesian analysis (Liao et al., 2020; Ju et al., 2023). Further work studies GNN generalization and transfer (Maskey et al., 2022; 2025; Levie, 2023; Vasileiou et al., 2025a; Rauchwerger et al., 2025); see also the survey of Vasileiou et al. (2025b). Our contribution differs by focusing on invariant estimation across changing dimensions, with explicit dimension-uniform rates and phase transitions driven by stable feature complexity.

Source conditions and optimal rates. Classical analyses of regularized least squares derive optimal statistical rates under source and capacity conditions (Caponnetto & De Vito, 2007). More recent work establishes exact and minimax rates for kernel ridge regression under source conditions in large dimensions (Zhang et al., 2025). These results concern fixed domains or joint asymptotics of sample size and ambient dimension. We instead study transfer from one object size to all larger sizes and determine when the resulting rates can be uniform over both the training and deployment sizes.

B FUNCTION-LEVEL SYMMETRY, STATISTICAL LEARNING, AND TRANSFER

This appendix formalizes the function-level framework of Section 3. It explains why unlabeled set and graph laws yield invariant density and regression targets, records representative losses under the unified observation notation, and defines exact transfer through coherent linear isometries. It then develops the stable invariant function spaces used in the rate calculations.

B.1 GROUP ACTIONS ON DATA AND FUNCTIONS

A group G acts on a measurable space $\mathcal{X}$ if

$$e \cdot x = x, \quad g \cdot (h \cdot x) = (gh) \cdot x,$$

for every $g, h \in G$ and $x \in \mathcal{X}$. This action induces a linear action on real-valued functions $f : \mathcal{X} \rightarrow \mathbb{R}$ by pullback:

$$(g \cdot f)(x) := f(g^{-1} \cdot x).$$

A function is G -invariant when $g \cdot f = f$ for every $g \in G$, equivalently,

$$f(g \cdot x) = f(x) \quad \text{for every } g \in G, x \in \mathcal{X}.$$

For sets and graphs of size d , the group is $G = S_d$, acting by relabeling elements or nodes as specified in Section 3.

A probability distribution μ on $\mathcal{X}$ is G -invariant if

$$g \cdot X \stackrel{\text{law}}{=} X \quad \text{for every } g \in G.$$

Equivalently, $\mu(g^{-1}A) = \mu(A)$ for all measurable sets $A \subseteq \mathcal{X}$.

Density targets. Suppose μ admits a density f^* with respect to a G -invariant reference measure λ . For every measurable A ,

$$\int_A f^* d\lambda = \mu(A) = \mu(g^{-1}A) = \int_A f^*(g^{-1} \cdot x) d\lambda(x).$$

Hence $f^*(g \cdot x) = f^*(x)$ for λ -almost every x . Thus an unlabeled set or graph distribution has an invariant density whenever its density is expressed relative to an invariant reference measure.

Regression targets. Let ρ be a joint distribution on $\mathcal{X} \times \mathbb{R}$ satisfying

$$(g \cdot X, Y) \stackrel{\text{law}}{=} (X, Y).$$

Then the conditional law of Y given $X = x$ agrees with that given $X = g \cdot x$, up to the usual almost-sure choice of a conditional version. Consequently, the regression function

$$f^*(x) = \mathbb{E}[Y \mid X = x]$$

admits a G -invariant version.

Invariant risk minimization. Let $\mathcal{F}$ be a function class closed under the induced G -action. A population risk $L : \mathcal{F} \rightarrow \mathbb{R}$ is G -invariant if $L(g \cdot f) = L(f)$. If the minimizer is unique, it is necessarily invariant. If the minimizer is not unique, $\mathcal{F}$ is convex, and L is convex, then group averaging produces an invariant minimizer:

$$\bar{f}(x) := \int_G (g \cdot f)(x) d\eta_G(g),$$

where η_G is the uniform measure for finite groups and Haar probability measure for compact groups. Indeed, Jensen's inequality and invariance give

$$L(\bar{f}) \leq \int_G L(g \cdot f) d\eta_G(g) = L(f).$$

This argument takes place entirely in function space and requires no action on a parameterization. Accordingly, uniqueness means uniqueness as an L^2 function, up to almost-everywhere equality; it is compatible with many parameter values representing that same function.

B.2 EXAMPLES OF STATISTICAL LEARNING OBJECTIVES

Recall the unified observation $Z_i = X_i$ for unsupervised learning and $Z_i = (X_i, Y_i)$ for supervised learning. In either case, $\hat{L}_{n,d}(f) = n^{-1} \sum_{i=1}^n \ell_d(Z_i, f)$ estimates $L_d(f) = \mathbb{E}[\ell_d(Z, f)]$. The formulation includes the following standard tasks.

Unsupervised statistical learning: density estimation. Let $\mathcal{F}_d \subseteq L^2(\lambda_d)$ be a class of nonnegative functions satisfying $\int f d\lambda_d = 1$. Since $\mu_d = f_d^* \lambda_d$, the negative log-likelihood loss

$$\ell_d(x, f) = -\log f(x)$$

has population risk

$$L_d(f) = \mathbb{E}_{X \sim \mu_d}[-\log f(X)].$$

This risk is uniquely minimized by f_d^* , up to λ_d -almost-everywhere equality, whenever $f_d^* \in \mathcal{F}_d$ and the risk is finite. Restricting to $\mathcal{F}_d^{\text{inv}}$ gives invariant maximum likelihood directly at the function level.

Supervised statistical learning: regression. For real-valued labels, squared loss gives

$$\ell_d(x, y, f) = (y - f(x))^2, \quad f_d^*(x) = \mathbb{E}[Y | X = x].$$

More generally, any proper supervised loss may be used when its population minimizer is the desired conditional target. If relabeling X does not change the joint law of (X, Y) , the target is invariant.

Score matching and denoising. The scalar formulation is not tied to likelihood: proper scoring rules, density-ratio objectives, and variational losses also apply whenever their population minimizer identifies f_d^* . Score matching and denoising instead estimate a vector-valued field, such as $\nabla \log f_d^*$, in $L^2(\lambda_d; \mathbb{R}^{m_d})$, where $m_d = rd$ for sets and $m_d = rd^2$ for graphs. Invariance of the density makes its score permutation equivariant. Replacing invariant scalar spaces by the corresponding equivariant vector spaces leaves the transfer, truncation, and estimation arguments unchanged, provided the same representation-stability and source-decay conditions hold.

B.3 EXACT FUNCTIONAL TRANSFER ACROSS SIZES

For density estimation let $\mathcal{H}_d = L^2(\lambda_d)$, and for regression let $\mathcal{H}_d = L^2(\mu_d)$. In either case, let $\mathcal{H}_d^{\text{inv}}$ be its invariant subspace, with inner product $\langle \cdot, \cdot \rangle_d$ and norm $\| \cdot \|_d$. An exact transfer system is a collection of linear maps

$$T_{d \rightarrow D} : \mathcal{H}_d^{\text{inv}} \longrightarrow \mathcal{H}_D^{\text{inv}}, \quad d \leq D,$$

satisfying the following three properties:

- (i) *coherence*: $T_{d \rightarrow d} = \text{Id}$ and $T_{D \rightarrow E} T_{d \rightarrow D} = T_{d \rightarrow E}$;
- (ii) *isometry*: $\langle T_{d \rightarrow D} f, T_{d \rightarrow D} h \rangle_D = \langle f, h \rangle_d$;
- (iii) *target compatibility*: $T_{d \rightarrow D} f_d^* = f_D^*$.

The isometry identity implies

$$\|T_{d \rightarrow D} f - T_{d \rightarrow D} h\|_D = \|f - h\|_d.$$

The map need not reach every function at the larger size; it only identifies the size-stable subspace carrying the shared rule.

These assumptions express the structure needed to learn at one size and make a statistical claim at another. Without target compatibility, the unseen target f_D^* is not identifiable from size- d observations. Without norm control, transfer could arbitrarily amplify a small source-size error. Isometry rules out that artifact and makes errors comparable across sizes. For score matching or denoising, the same definition applies to the corresponding equivariant vector-valued spaces.

Stable invariant features. Suppose $\{\psi_{d,j}\}_{j \geq 1}$ and $\{\psi_{D,j}\}_{j \geq 1}$ are corresponding orthonormal invariant features. Define

$$T_{d \rightarrow D} \psi_{d,j} = \psi_{D,j}$$

and extend linearly and continuously. Then, for every square-summable sequence $(a_j)_j$,

$$T_{d \rightarrow D} \left(\sum_j a_j \psi_{d,j} \right) = \sum_j a_j \psi_{D,j}, \quad \left\| T_{d \rightarrow D} \sum_j a_j \psi_{d,j} \right\|_D^2 = \sum_j a_j^2.$$

Thus transfer preserves the learned rule and its error exactly. This construction underlies the stable polynomial and Hermite classes below. It is also the function-space abstraction of weight sharing in Deep Sets and GNNs: the computation is instantiated on a new number of elements or nodes, but the rule itself is unchanged (Zaheer et al., 2017; Xu et al., 2019; Levin & Díaz, 2024).

Common ambient objects. Exact transfer also arises when all finite-size functions are realizations of a single rule on a common limit object. Mean embeddings and normalized empirical measures provide such objects for sets. Graphons and normalized motif functionals provide them for dense graphs (Ruiz et al., 2020; Diaconis & Janson, 2008). Appendix C gives explicit set and graph constructions, including replication, graph blow-ups, and shared Deep Sets and GNN rules.

B.4 INVARIANT POLYNOMIAL FUNCTION SPACES

A central example is a finite-dimensional space of polynomial functions. Let V_d be a finite-dimensional vector space on which G_d acts linearly, and let $\mathcal{P}_{d,k}$ be the space of homogeneous degree- k polynomial functions on V_d . The induced action on functions is

$$(g \cdot f)(x) := f(g^{-1} \cdot x).$$

The invariant homogeneous polynomial space is

$$\mathcal{P}_{d,k}^{G_d} := \{f \in \mathcal{P}_{d,k} : g \cdot f = f \text{ for all } g \in G_d\}.$$

For density estimation, completeness follows from the Hermite basis of $L^2(\lambda_d)$. For regression, we assume the standard polynomial-completeness condition in $L^2(\mu_d)$, which holds, for example, under suitable moment determinacy or exponential-moment conditions. Averaging polynomial approximants over S_d then proves density of invariant polynomials in $\mathcal{H}_d^{\text{inv}}$.

For finite groups, the dimension of the invariant subspace can be expressed using characters. If $\varrho_{d,k}$ is the representation of G_d on $\mathcal{P}_{d,k}$ (i.e., the collection of matrices corresponding to the action) and $\chi_{d,k}(g) = \text{tr}(\varrho_{d,k}(g))$ is its character, then

$$\dim \mathcal{P}_{d,k}^{G_d} = \frac{1}{|G_d|} \sum_{g \in G_d} \chi_{d,k}(g).$$

Equivalently, if P_{G_d} denotes the group-averaging projection

$$P_{G_d} f = \frac{1}{|G_d|} \sum_{g \in G_d} g \cdot f,$$

then

$$\dim \mathcal{P}_{d,k}^{G_d} = \text{tr}(P_{G_d}).$$

For compact groups, the finite average is replaced by Haar integration.

Representation stability is the phenomenon that these invariant function-space dimensions stabilize as d grows. This stabilization is the algebraic mechanism behind dimension-uniform learning in our setting.

B.5 SET-VALUED DATA AND MULTISYMMETRIC POLYNOMIALS

Consider set-valued data with d elements in $\mathbb{R}^r$. We represent the set as

$$X = (x_1, \dots, x_d) \in (\mathbb{R}^r)^d, \quad x_i = (x_{i,1}, \dots, x_{i,r}) \in \mathbb{R}^r.$$

The symmetric group S_d acts by permuting the d elements:

$$\pi \cdot X = (x_{\pi^{-1}(1)}, \dots, x_{\pi^{-1}(d)}).$$

The corresponding invariant polynomials are called multisymmetric polynomials: they are polynomials in the dr variables $x_{i,a}$ that are invariant under permutations of the block index i .

Let $\mathcal{P}_{d,r,k}^{S_d}$ denote the space of homogeneous degree k multisymmetric polynomials. A monomial is described by a collection of multi-indices

$$\alpha_i = (\alpha_{i,1}, \dots, \alpha_{i,r}) \in \mathbb{N}^r, \quad 1 \leq i \leq d,$$

with total degree $\sum_i |\alpha_i| = k$. The group S_d permutes the multi-indices α_i . Therefore an invariant monomial orbit is determined by a multiset of nonzero multi-indices

$$\{\alpha^{(1)}, \dots, \alpha^{(q)}\}, \quad \alpha^{(j)} \in \mathbb{N}^r \setminus \{0\}, \quad \sum_{j=1}^q |\alpha^{(j)}| = k.$$

For $d \geq k$, every such multiset can be realized using at most k nonzero blocks, so the dimension stabilizes in d .

For each $m \geq 1$, the number of nonzero block types of degree m is

$$a_r(m) = \#\{\alpha \in \mathbb{N}^r : |\alpha| = m\} = \binom{r+m-1}{m}.$$

Hence the stable dimension of homogeneous degree k invariants is

$$N_r(k) := \dim \mathcal{P}_{d,r,k}^{S_d} = [t^k] \prod_{m \geq 1} (1 - t^m)^{-a_r(m)} \quad \text{for } d \geq k,$$

where $[t^k]F(t)$ denotes the coefficient of t^k in the formal power series $F(t)$. Equivalently,

$$N_r(k) = [t^k] \prod_{m \geq 1} (1 - t^m)^{-\binom{r+m-1}{m}}.$$

For $r = 1$, this reduces to the ordinary partition number:

$$N_1(k) = p(k).$$

Thus the invariant degree- k dimension is independent of d once $d \geq k$, whereas the full homogeneous degree- k polynomial space in dr variables has dimension

$$\binom{dr+k-1}{k},$$

which grows polynomially in d of order d^k for fixed k .

The asymptotic behavior of $N_r(k)$ for fixed r and large k is subexponential:

$$\log N_r(k) \sim C_r k^{\frac{r}{r+1}},$$

where

$$C_r = \frac{r+1}{r} (r\zeta(r+1))^{\frac{1}{r+1}},$$

and $\zeta(s) := \sum_{n=1}^{\infty} n^{-s}$ denotes the Riemann zeta function. Thus

$$N_r(k) = \exp \left(C_r k^{\frac{r}{r+1}} + o\left(k^{\frac{r}{r+1}}\right) \right).$$

For example, $r = 1$ recovers the Hardy–Ramanujan exponent

$$\log p(k) \sim \pi \sqrt{\frac{2k}{3}}.$$

For $r = 2$,

$$\log N_2(k) \sim \frac{3}{2} (2\zeta(3))^{1/3} k^{2/3},$$

and for $r = 3$,

$$\log N_3(k) \sim \frac{4}{3} (3\zeta(4))^{1/4} k^{3/4}.$$

These formulas show that for fixed element dimension r and polynomial degree k , the invariant polynomial function-space dimension stabilizes as the set size d grows. This is the representation-stability mechanism underlying the dimension-independent rates for set-valued data.

B.6 HERMITE FEATURES FOR SET-VALUED DATA

For data on $(\mathbb{R}^r)^d$, a natural basis is given by multivariate Hermite polynomials relative to a Gaussian reference measure. Let H_α denote an orthonormal Hermite polynomial indexed by $\alpha \in \mathbb{N}^{dr}$. For block-structured data, we may write

$$H_{\alpha_1, \dots, \alpha_d}(X) = \prod_{i=1}^d H_{\alpha_i}(x_i), \quad \alpha_i \in \mathbb{N}^r.$$

The group S_d permutes the block indices α_i . Therefore invariant Hermite features are obtained by symmetrizing Hermite monomials over S_d . The dimension count for homogeneous degree k invariant Hermite features is the same as for multisymmetric polynomials:

$$\dim \mathcal{H}_{d,r,k}^{S_d} = N_r(k) \quad \text{for } d \geq k.$$

This construction gives a concrete hierarchy for invariant density and regression functions. For density estimation, let $\phi_{d,j}$ be invariant Hermite functions; for regression, let them be the corresponding invariant polynomials orthonormalized in $L^2(\mu_d)$. In either case, the scalar target admits an expansion

$$f_d^*(X) = \sum_j a_{d,j}^* \phi_{d,j}(X).$$

Truncation at degree k produces a finite-dimensional invariant function class whose dimension is

$$N_r(\leq k) = \sum_{q=0}^k N_r(q),$$

independent of d once $d \geq k$. For density estimation, the admissible class additionally enforces nonnegativity and unit integral.

If one instead studies score matching, an invariant log-density expansion $\log f_d^* = \sum_j b_{d,j}^* \phi_{d,j}$ gives the Lebesgue score

$$\nabla_X \log \frac{d\mu_d}{dx}(X) = -X + \sum_j b_{d,j}^* \nabla_X \phi_{d,j}(X).$$

Gradients of invariant scalar features are equivariant vector fields:

$$\nabla_X \phi_{d,j}(\pi \cdot X) = \pi \cdot \nabla_X \phi_{d,j}(X).$$

If the function coefficients have a controlled tail, for example,

$$\sum_{j>m} (a_{d,j}^*)^2 \leq b_m^2,$$

then truncating to m invariant Hermite functions incurs squared L^2 approximation error at most b_m^2 . This is a concrete instance of the approximation tail $B(k)$ in Theorem 4.1.

B.7 ATTRIBUTED GRAPHS

An attributed graph with d nodes is represented, as in the main text, by

$$X \in \mathcal{X}_d = \mathbb{R}^{r \times d \times d}.$$

The channel index $a \in \{1, \dots, r\}$ records edge and/or node features; node features may, for example, be placed on diagonal entries of designated channels. Node relabeling acts simultaneously on both node indices:

$$(\pi \cdot X)_{a,i,j} = X_{a,\pi^{-1}(i),\pi^{-1}(j)}.$$

A homogeneous degree- k polynomial function is a linear combination of monomials

$$\prod_{a=1}^r \prod_{i,j=1}^d X_{a,i,j}^{m_{a,i,j}}, \quad \sum_{a,i,j} m_{a,i,j} = k.$$

Two such monomials belong to the same S_d -orbit precisely when their node indices differ only by a relabeling. The orbits are therefore indexed by unlabeled channel-attributed directed multigraph patterns of total degree k , with diagonal factors representing vertex decorations when node features are encoded on the diagonal.

Let $\mathcal{R}_{d,r,k}^{S_d}$ be the corresponding space of invariant homogeneous polynomial functions. Any degree- k monomial involves at most $2k$ distinct node indices, so

$$\dim \mathcal{R}_{d,r,k}^{S_d} = G_r(k) \quad \text{for every } d \geq 2k,$$

where $G_r(k)$ is the number of admissible unlabeled channel-attributed multigraph patterns of degree k . Thus the function-space dimension stabilizes with respect to the number of nodes.

The same dimension can be written by character theory. With

$$W_d = \mathbb{R}^{r \times d \times d},$$

we have

$$\mathcal{R}_{d,r,k}^{S_d} = \left(\text{Sym}^k(W_d^*) \right)^{S_d}, \quad \dim \mathcal{R}_{d,r,k}^{S_d} = \frac{1}{d!} \sum_{\pi \in S_d} \chi_{\text{Sym}^k(W_d^*)}(\pi).$$

Equivalently, if ϱ_d is the representation of S_d on W_d^* , its Molien series is

$$\sum_{k \geq 0} \dim \mathcal{R}_{d,r,k}^{S_d} t^k = \frac{1}{d!} \sum_{\pi \in S_d} \frac{1}{\det(I - t\varrho_d(\pi))}.$$

The asymptotic dependence on k separates node-only and edge-attributed systems. If all active channels are supported on the diagonal, the stable dimension reduces to the multisymmetric count

$$N_r(k) = [t^k] \prod_{m \geq 1} (1 - t^m)^{-\binom{r+m-1}{m}},$$

and therefore

$$N_r(k) = \exp \left(C_r k^{\frac{r}{r+1}} + o \left(k^{\frac{r}{r+1}} \right) \right).$$

In contrast, if at least one channel contains edge variables, graph-pattern complexity gives

$$G_r(k) = \exp(k \log k + O(k \log \log k)), \quad k \rightarrow \infty.$$

Thus attributed-graph invariant function spaces stabilize in the number of nodes d , while their growth with interaction order k is larger than for set-valued data.

C CONCRETE EXACT TRANSFERS FOR SETS AND GRAPHS

This appendix makes the transfer assumption concrete for the two domains studied in the paper. The operator $T_{d \rightarrow D}$ does not match arbitrary functions across dimensions; it instantiates the same invariant rule at a new set or graph size. Shared Deep Sets, GNNs, empirical measures, and graphons motivate coherence and exact target compatibility. The stable feature constructions additionally realize the linear isometry required by the theory after normalization in the relevant L^2 spaces.

C.1 SETS

A set of d vectors is represented by $X = (x_1, \dots, x_d) \in \mathcal{X}_d = (\mathbb{R}^r)^d \simeq \mathbb{R}^{r \times d}$. Because the ordering carries no information, S_d acts by $\pi \cdot X = (x_{\pi^{-1}(1)}, \dots, x_{\pi^{-1}(d)})$. A set distribution is therefore represented by an exchangeable law, and both its density and its regression function are invariant by Appendix B.1.

Shared Deep Sets rules. Let $\phi : \mathbb{R}^r \rightarrow \mathbb{R}^m$ be an elementwise feature map and define the normalized set summary

$$m_d^\phi(X) = \frac{1}{d} \sum_{i=1}^d \phi(x_i).$$

For any readout $F : \mathbb{R}^m \rightarrow \mathbb{R}$, the rule

$$f_d(X) = F \left(m_d^\phi(X) \right)$$

is invariant for every d . Its transfer to size D is obtained by applying the same ϕ and F :

$$(T_{d \rightarrow D} f_d)(X_1, \dots, X_D) = F \left(\frac{1}{D} \sum_{i=1}^D \phi(X_i) \right).$$

This is precisely the size-shared computation of a mean-pooled Deep Sets model. Sum-pooled models admit the same construction when the task uses an extensive, rather than normalized, quantity. Exact target compatibility means that the population task is generated by this same rule at every size.

Replication and empirical measures. If $D = qd$, repeat each element q times:

$$R_{d \rightarrow D}(x_1, \dots, x_d) = (\underbrace{x_1, \dots, x_1}_q, \dots, \underbrace{x_d, \dots, x_d}_q).$$

The normalized summary is unchanged,

$$m_D^\phi(R_{d \rightarrow D}X) = m_d^\phi(X).$$

Equivalently, replication leaves the empirical measure $d^{-1} \sum_i \delta_{x_i}$ unchanged. Hence every function of the empirical measure, including normalized moments and mean-pooled Deep Sets rules, transfers exactly. Zero padding gives another exact construction for sum-based rules whenever the added null element satisfies $\phi(0) = 0$.

Stable feature expansions. Let $\psi_{d,j}$ be normalized multisymmetric or invariant Hermite functions, paired across sizes by their common orbit type. For

$$f_d = \sum_j a_j \psi_{d,j},$$

define

$$T_{d \rightarrow D} f_d = \sum_j a_j \psi_{D,j}.$$

If the paired features are orthonormal in their respective task spaces, Parseval's identity gives

$$\|T_{d \rightarrow D} f_d\|_D^2 = \sum_j a_j^2 = \|f_d\|_d^2.$$

Thus $T_{d \rightarrow D}$ is a linear isometry. A shared expansion may represent either a regression function or a density; in the latter case, the admissible coefficient class is restricted so that the expansion remains nonnegative and integrates to one. This provides the explicit transfer used by the set function classes in Appendix B.5.

C.2 ATTRIBUTED GRAPHS

An attributed graph with d nodes is represented by $X \in \mathcal{X}_d = \mathbb{R}^{r \times d \times d}$, and S_d acts by simultaneous relabeling:

$$(\pi \cdot X)_{a,i,j} = X_{a,\pi^{-1}(i),\pi^{-1}(j)}.$$

Consequently, graph-level densities and regression functions depend only on the isomorphism class of the attributed graph.

Normalized motif rules and graph blow-ups. For a fixed attributed pattern H , let $t_H(X)$ denote its normalized homomorphism density in X . These functions are invariant and are indexed by H , not by the number of nodes. If $X^{(q)}$ is the q -fold blow-up obtained by replacing each node by q twins and each edge by the corresponding complete bipartite block, then

$$t_H(X^{(q)}) = t_H(X).$$

Thus graph blow-ups preserve every rule of the form

$$f_d(X) = F(t_{H_1}(X), \dots, t_{H_m}(X))$$

exactly. A step-function graphon gives the same interpretation: refining its finite partition changes the graph size but not the common graphon or its normalized motif functionals.

Shared GNN rules. A message-passing GNN applies the same local aggregation and update functions at every node and every size, followed by a permutation-invariant graph readout. Holding these functions fixed while changing d therefore defines a map from the size- d function to its size- D realization. This is the architecture-level instance of $T_{d \rightarrow D}$. The transfer is exact when the target family is generated by the same shared GNN rule. Mean aggregation and normalized graph readouts prevent the numerical scale of the rule from growing merely with the number of nodes.

Stable graph feature expansions. Let $\psi_{d,H}$ be normalized invariant functions indexed by unlabeled attributed multigraph patterns H . Transfer pairs the same pattern across sizes:

$$T_{d \rightarrow D} \psi_{d,H} = \psi_{D,H}.$$

Extending this identification linearly gives

$$T_{d \rightarrow D} \left(\sum_H a_H \psi_{d,H} \right) = \sum_H a_H \psi_{D,H}.$$

After orthonormalization in the task norms, this map is an isometry and satisfies the coherence identity. The degree- k truncations stabilize once $d \geq 2k$, as shown in Appendix B.7. These expansions cover invariant graph regression and density functions without privileging a particular learning loss; for densities, admissibility again enforces nonnegativity and normalization.

C.3 WHAT THE EXACT-TRANSFER ASSUMPTION ASSERTS

Across all of these examples, exact transfer makes two statements. First, the semantic rule is shared across sizes: the same normalized summary, architecture, motif functional, or invariant expansion is used at the training and target sizes. Second, the task family is compatible with that rule:

$$T_{d \rightarrow D} f_d^* = f_D^*.$$

This does not claim that arbitrary set or graph tasks are related. It identifies the broad class of size-consistent problems for which learning at one size can be statistically informative at another. The isometry assumption then ensures that this semantic identification does not alter the scale of estimation error. Together, these conditions isolate the statistical question studied in the paper from a separate cross-size modeling mismatch.

D PROOF OF THEOREM 4.1

This section proves the dimension-free upper bounds and derives all rate regimes. The proof has two steps. First, exact isometric transfer converts a fixed-size estimation guarantee into a simultaneous guarantee at every larger size. Second, representation growth is balanced against the source tail.

Exact transfer is defined in Section 3 and formalized in Appendix B.3; representation growth $q_X(k)$ and the source envelope $B(k)$ are defined in Section 4. The loss assumptions in the main theorem are used below to derive the finite-dimensional estimation bound. The minimax theorem additionally uses the local nondegeneracy conditions equation E.2–equation E.3.

D.1 FINITE-DIMENSIONAL ESTIMATION AND TRANSFER

Fix a domain type $X \in \{\text{set}, \text{graph}\}$, a training size d , and an interaction order k . Let $f_{d,k}^\circ$ and $\hat{f}_{n,d,k}^{\text{inv}}$ minimize the population and empirical risks over $\mathcal{F}_{d,k}$, respectively. Uniform calibration gives $\|f_{d,k}^\circ - f_d^*\|_d^2 \leq C_{\text{app}} B_d(k)$. Apply the strong-convexity and score-concentration calculation of Appendix F in orthonormal coordinates on $\mathcal{V}_{d,k}$. Since those coordinates identify Euclidean distance with $\|\cdot\|_d$, with probability at least $1 - \delta$,

$$\|\hat{f}_{n,d,k}^{\text{inv}} - f_{d,k}^\circ\|_d^2 \leq C \frac{\sigma^2 \dim(\mathcal{V}_{d,k}) + \log(1/\delta)}{\alpha^2 n}.$$

The squared triangle inequality and the calibration bound therefore give

$$\|\hat{f}_{n,d,k}^{\text{inv}} - f_d^*\|_d^2 \leq C_0 \left(B_d(k) + \frac{\dim(\mathcal{V}_{d,k}) + \log(1/\delta)}{n} \right). \quad (\text{D.1})$$

Here C_0 depends on the uniform calibration, curvature, and score constants, but not on n, d, D , or k . Thus equation D.1 is a consequence of the primitive loss conditions, rather than an assumed rate.

The two quantities on the right are uniform in size: by definition, $B_d(k) \leq B(k)$ and $\dim(\mathcal{V}_{d,k}) \leq q_X(k)$. Moreover, exact target transfer and isometry give, simultaneously for every $D \geq d$,

$$\begin{aligned} \|T_{d \rightarrow D} \hat{f}_{n,d,k}^{\text{inv}} - f_D^*\|_D^2 &= \|T_{d \rightarrow D} (\hat{f}_{n,d,k}^{\text{inv}} - f_d^*)\|_D^2 \\ &= \|\hat{f}_{n,d,k}^{\text{inv}} - f_d^*\|_d^2. \end{aligned} \quad (\text{D.2})$$

No union bound over D is needed: all target-size errors equal the same source-size error on the event in equation D.1. Taking the supremum in equation D.2 therefore yields

$$\sup_{D \geq d} \|T_{d \rightarrow D} \hat{f}_{n,d,k}^{\text{inv}} - f_D^*\|_D^2 \leq C_0 \left(B(k) + \frac{q_{\mathbf{x}}(k) + \log(1/\delta)}{n} \right). \quad (\text{D.3})$$

This is the key dimension-free oracle inequality. Object sizes d and D have disappeared; only interaction order, representation growth, and target regularity remain.

D.2 SET RATES

Write $\beta = r/(r+1)$. The stable multisymmetric count satisfies

$$\log q_{\text{set}}(k) = C_r k^\beta + o(k^\beta), \quad C_r = \frac{r+1}{r} (r\zeta(r+1))^{1/(r+1)}. \quad (\text{D.4})$$

We optimize the two leading terms in equation D.3; for a fixed confidence level, $\log(1/\delta)/n$ is smaller than all rates below.

Polynomial source decay. If $B(k) \lesssim k^{-\tau}$, balancing the two terms in equation D.3 gives $C_r k_n^\beta + \tau \log k_n = \log n + o(\log n)$, hence $k_n = (\log n / C_r)^{1/\beta} (1 + o(1))$ and

$$\sup_{D \geq d} \|T_{d \rightarrow D} \hat{f}_{n,d,k_n}^{\text{inv}} - f_D^*\|_D^2 \lesssim (\log n)^{-\tau/\beta} = (\log n)^{-\tau(r+1)/r}. \quad (\text{D.5})$$

Exponential source decay. If $B(k) \lesssim e^{-ak^\nu}$, the balancing equation is

$$ak_n^\nu + C_r k_n^\beta + o(k_n^\beta) = \log n. \quad (\text{D.6})$$

Solving it in the three regimes gives the compact summary

$$B(k_n) + \frac{q_{\text{set}}(k_n)}{n} \asymp \begin{cases} n^{-1+o(1)}, & \nu > \beta, \\ n^{-a/(a+C_r)+o(1)}, & \nu = \beta, \\ \exp[-aC_r^{-\nu/\beta} (\log n)^{\nu/\beta} (1+o(1))], & 0 < \nu < \beta. \end{cases} \quad (\text{D.7})$$

The first line is the fast set regime in Table 2; the other two lines give the critical and subcritical interpolation.

D.3 GRAPH RATES

For every fixed feature dimension r , the stable graph-pattern count satisfies

$$\log q_{\text{graph}}(k) = k \log k + O_r(k \log \log k) = k \log k (1 + o(1)). \quad (\text{D.8})$$

Thus r affects only the lower-order remainder and does not enter the first-order rates.

Polynomial source decay. For $B(k) \lesssim k^{-\tau}$, the crossing equation gives $k_n = (\log n / \log \log n)(1 + o(1))$, and therefore

$$\sup_{D \geq d} \|T_{d \rightarrow D} \hat{f}_{n,d,k_n}^{\text{inv}} - f_D^*\|_D^2 \lesssim \left(\frac{\log n}{\log \log n} \right)^{-\tau}. \quad (\text{D.9})$$

Exponential source decay. For $B(k) \lesssim e^{-ak^\nu}$, the balancing equation is

$$ak_n^\nu + k_n \log k_n + O_r(k_n \log \log k_n) = \log n. \quad (\text{D.10})$$

Solving it gives

$$B(k_n) + \frac{q_{\text{graph}}(k_n)}{n} \asymp \begin{cases} n^{-1+o(1)}, & \nu > 1, \\ \exp\left[-a \left(\frac{\log n}{\log \log n} \right)^\nu (1+o(1))\right], & 0 < \nu \leq 1. \end{cases} \quad (\text{D.11})$$

The second line is critical at $\nu = 1$ and subcritical for $0 < \nu < 1$. Equations equation D.5, equation D.7, equation D.9, and equation D.11 prove Theorem 4.1.

E PROOF OF THEOREM 4.2

We now show that the preceding rates cannot be improved. The lower bound is stated under the nondegeneracy condition in the theorem, which requires the invariant coordinates counted by representation stability to correspond to statistically distinguishable local perturbations. This is the standard condition that prevents the loss or data distribution from collapsing nominally different invariant directions.

E.1 MINIMAX RISK AND LOCAL INVARIANT SUBMODELS

Let $\mathfrak{F}_X(B)$ be the class of size-consistent target families on domain X satisfying exact transfer and the source envelope B . An estimator observes n samples at size d and returns functions $\hat{f}_D$ for $D \geq d$. Define its worst-case risk by

$$\mathfrak{R}_n(X, B) := \inf_{\hat{f}} \sup_{d \geq 1} \sup_{f^* \in \mathfrak{F}_X(B)} \mathbb{E} \left[\sup_{D \geq d} \|\hat{f}_D - f_D^*\|_D^2 \right]. \quad (\text{E.1})$$

Minimax nondegeneracy. For every interaction order k , take a size d_k in the stable range: $d_k \geq k$ for sets and $d_k \geq 2k$ for graphs. The nondegeneracy assumption provides $m_k \geq c_0 q_X(k)$ orthonormal invariant directions $\psi_1, \dots, \psi_{m_k}$, paired coherently across larger sizes, and an admissible baseline f_0 such that the hypercube

$$f_\omega = f_0 + u \sum_{j=1}^{m_k} \omega_j \psi_j, \quad \omega \in \{-1, 1\}^{m_k}, \quad (\text{E.2})$$

belongs to the source class whenever $m_k u^2 \leq c_1 B(k)$. It also requires the neighboring statistical experiments to satisfy

$$\text{KL}(P_\omega^{\otimes n}, P_{\omega^{(j)}}^{\otimes n}) \leq c_2 n u^2, \quad (\text{E.3})$$

where $\omega^{(j)}$ differs from ω only in coordinate j . For regression, one obtains equation E.3 by adding Gaussian noise to $Y = f_\omega(X)$. For density estimation, it follows from a bounded local density or exponential-family perturbation around an invariant density bounded away from zero. These constructions also ensure nonnegativity and normalization of every density in equation E.2.

E.2 ASSOUD LOWER BOUND

Set

$$u_k^2 = c_3 \min \left\{ \frac{B(k)}{m_k}, \frac{1}{n} \right\},$$

with c_3 sufficiently small. Then the hypercube is contained in the source class, and the KL divergence between neighboring vertices is bounded by a universal constant. Since orthonormality gives $\|f_\omega - f_{\omega^{(j)}}\|_{d_k}^2 = 4u_k^2$, Assouad's lemma yields

$$\inf_{\hat{f}} \sup_{\omega \in \{-1, 1\}^{m_k}} \mathbb{E}_\omega \|\hat{f} - f_\omega\|_{d_k}^2 \geq c_4 m_k u_k^2 \geq c_5 \min \left\{ B(k), \frac{q_X(k)}{n} \right\}. \quad (\text{E.4})$$

The size-generalization loss contains the choice $D = d_k$, so its supremum is at least the in-domain loss in equation E.4. Consequently,

$$\mathfrak{R}_n(X, B) \geq c_5 \sup_{k \geq 1} \min \left\{ B(k), \frac{q_X(k)}{n} \right\}. \quad (\text{E.5})$$

This observation is important: transfer cannot remove a lower bound already present at the training size.

Because $B(k)$ decreases and $q_X(k)$ increases, the supremum in equation E.5 is attained, up to constants and lower-order factors, where $B(k) \asymp q_X(k)/n$. Under the regular source envelopes considered here,

$$\sup_k \min \left\{ B(k), \frac{q_X(k)}{n} \right\} \asymp \inf_k \left\{ B(k) + \frac{q_X(k)}{n} \right\}. \quad (\text{E.6})$$

The right-hand side is precisely the quantity controlled by the upper oracle inequality equation D.3.

E.3 EVALUATION OF THE LOWER BOUND

For polynomial source decay, substituting the set and graph representation asymptotics into the crossing condition gives, respectively,

$$(\log n)^{-\tau(r+1)/r} \quad \text{and} \quad \left(\frac{\log n}{\log \log n} \right)^{-\tau}. \quad (\text{E.7})$$

For exponential source decay, the same crossing equations give exactly the three set regimes in equation D.7 and the two graph regimes in equation D.11.

These lower bounds match the upper bounds in Appendix D, up to constants and lower-order factors. They apply to both regression and density estimation through the local submodels above, and they already hold at a stable training size. Therefore no estimator can uniformly improve any set or graph rate in Table 2, completing the proof of Theorem 4.2.

F AUXILIARY FINITE-DIMENSIONAL ESTIMATION BOUND

For completeness, we derive the finite-dimensional estimation term in equation D.1 from restricted curvature and score concentration. Empirical optimality and strong convexity reduce estimation error to the empirical gradient at the best population fit in the finite model; no statistical rate is assumed.

F.1 RESTRICTED GRADIENTS ON THE INVARIANT MODEL

Fix $d \in \mathbb{N}$. Let $\Theta_d^{G_d}$ be a finite-dimensional invariant parameter space and let $\theta_d^* \in \Theta_d^{G_d}$ denote its population-risk minimizer. In this section, θ_d^* is the best fit within the finite model (the analogue of $f_{d,k}^\circ$), and need not equal the unrestricted target. We write

$$p_d^G = \dim(\Theta_d^{G_d}).$$

For simplicity of notation, we identify $\Theta_d^{G_d}$ locally with a Euclidean space of dimension p_d^G . Equivalently, if $\Theta_d^{G_d}$ is a smooth model, the gradient below is understood in any normalized local coordinate chart around θ_d^* . Thus, $\nabla_G \hat{L}_{n,d}(\theta_d^*)$ denotes the gradient of the empirical risk restricted to the invariant model.

We work on the restricted-curvature event on which the empirical risk

$$\hat{L}_{n,d}(\theta) = \frac{1}{n} \sum_{i=1}^n \ell_d(Z_i, \theta)$$

is α -strongly convex on $\Theta_d^{G_d}$, with α independent of d . This is a primitive curvature property, not an estimation-rate assumption. For squared-loss regression in an orthonormal feature basis, the Hessian is the empirical feature Gram matrix. If the whitened feature vector is uniformly sub-Gaussian, standard covariance concentration gives this event with probability at least $1 - \delta$ once $n \gtrsim p_d^G + \log(1/\delta)$. For a regular likelihood or smooth proper loss, the same conclusion follows from a uniformly positive population Hessian and concentration of the restricted empirical Hessian. Pointwise strong convexity of the loss is a stronger sufficient condition, but is not required.

We will use the following standard concentration bound for the restricted empirical gradient.

Lemma F.1 (Restricted score concentration). *Suppose the restricted score at θ_d^* is sub-Gaussian with variance proxy at most σ^2 . That is, writing*

$$Y_i := \nabla_G \ell_d(Z_i, \theta_d^*) \in \mathbb{R}^{p_d^G},$$

we have $\mathbb{E}Y_i = 0$ and, for every unit vector $v \in \mathbb{R}^{p_d^G}$,

$$\mathbb{E} \exp(\lambda \langle v, Y_i \rangle) \leq \exp\left(\frac{\lambda^2 \sigma^2}{2}\right) \quad \forall \lambda \in \mathbb{R}.$$

Then there exists an absolute constant $C_0 > 0$ such that, for every $\delta \in (0, 1)$, with probability at least $1 - \delta$,

$$\left\| \nabla_G \hat{L}_{n,d}(\theta_d^*) \right\|_2^2 = \left\| \frac{1}{n} \sum_{i=1}^n Y_i \right\|_2^2 \leq C_0 \sigma^2 \frac{p_d^G + \log(1/\delta)}{n}.$$

Proof. Let $p = p_d^G$ and define

$$\bar{Y}_n := \frac{1}{n} \sum_{i=1}^n Y_i.$$

Since the Y_i 's are independent, mean-zero, and σ^2 -sub-Gaussian, the average $\bar{Y}_n$ is sub-Gaussian with variance proxy σ^2/n . Hence for every unit vector $v \in \mathbb{R}^p$,

$$\mathbb{E} \exp(\lambda \langle v, \bar{Y}_n \rangle) \leq \exp\left(\frac{\lambda^2 \sigma^2}{2n}\right).$$

Let $\mathcal{N}$ be a $1/2$ -net of the unit sphere $\mathbb{S}^{p-1}$ with $|\mathcal{N}| \leq 5^p$. For every $u \in \mathbb{R}^p$,

$$\|u\|_2 \leq 2 \max_{v \in \mathcal{N}} \langle v, u \rangle.$$

Therefore, for any $t > 0$,

$$\mathbb{P}(\|\bar{Y}_n\|_2 \geq 2t) \leq \mathbb{P}\left(\max_{v \in \mathcal{N}} \langle v, \bar{Y}_n \rangle \geq t\right).$$

By a union bound and the one-dimensional sub-Gaussian tail bound,

$$\mathbb{P}\left(\max_{v \in \mathcal{N}} \langle v, \bar{Y}_n \rangle \geq t\right) \leq |\mathcal{N}| \exp\left(-\frac{nt^2}{2\sigma^2}\right) \leq \exp\left(p \log 5 - \frac{nt^2}{2\sigma^2}\right).$$

Taking

$$t = \sigma \sqrt{\frac{2(p \log 5 + \log(1/\delta))}{n}},$$

we obtain, with probability at least $1 - \delta$,

$$\|\bar{Y}_n\|_2 \leq 2\sigma \sqrt{\frac{2(p \log 5 + \log(1/\delta))}{n}}.$$

Squaring both sides yields

$$\|\bar{Y}_n\|_2^2 \leq 8\sigma^2 \frac{p \log 5 + \log(1/\delta)}{n}.$$

Thus the claim holds with an absolute constant C_0 . □

F.2 PROOF OF THE ESTIMATION BOUND

Let

$$\hat{\theta}_{n,d}^G \in \arg \min_{\theta \in \Theta_d^{G_d}} \hat{L}_{n,d}(\theta).$$

Since $\theta_d^* \in \Theta_d^{G_d}$, empirical optimality gives

$$\hat{L}_{n,d}(\hat{\theta}_{n,d}^G) \leq \hat{L}_{n,d}(\theta_d^*).$$

On the intersection of the restricted-curvature event and the score-concentration event, $\hat{L}_{n,d}$ is α -strongly convex on $\Theta_d^{G_d}$, and hence

$$\hat{L}_{n,d}(\hat{\theta}_{n,d}^G) \geq \hat{L}_{n,d}(\theta_d^*) + \left\langle \nabla_G \hat{L}_{n,d}(\theta_d^*), \hat{\theta}_{n,d}^G - \theta_d^* \right\rangle + \frac{\alpha}{2} \left\| \hat{\theta}_{n,d}^G - \theta_d^* \right\|_2^2.$$

Combining the previous two displays gives

$$0 \geq \left\langle \nabla_G \hat{L}_{n,d}(\theta_d^*), \hat{\theta}_{n,d}^G - \theta_d^* \right\rangle + \frac{\alpha}{2} \left\| \hat{\theta}_{n,d}^G - \theta_d^* \right\|_2^2.$$

By Cauchy–Schwarz,

$$0 \geq - \left\| \nabla_G \hat{L}_{n,d}(\theta_d^*) \right\|_2 \left\| \hat{\theta}_{n,d}^G - \theta_d^* \right\|_2 + \frac{\alpha}{2} \left\| \hat{\theta}_{n,d}^G - \theta_d^* \right\|_2^2.$$

If $\hat{\theta}_{n,d}^G = \theta_d^*$, the desired bound is immediate. Otherwise, dividing by $\|\hat{\theta}_{n,d}^G - \theta_d^*\|_2$ gives

$$\left\| \hat{\theta}_{n,d}^G - \theta_d^* \right\|_2 \leq \frac{2}{\alpha} \left\| \nabla_G \hat{L}_{n,d}(\theta_d^*) \right\|_2.$$

Taking each event with failure probability $\delta/2$, squaring, applying Lemma F.1, and using a union bound, we obtain with probability at least $1 - \delta$,

$$\left\| \hat{\theta}_{n,d}^G - \theta_d^* \right\|_2^2 \leq \frac{4}{\alpha^2} \left\| \nabla_G \hat{L}_{n,d}(\theta_d^*) \right\|_2^2 \leq C \frac{\sigma^2 p_d^G + \log(1/\delta)}{\alpha^2 n},$$

for an absolute constant $C > 0$. This is the finite-dimensional estimation bound used in equation D.1. When the coordinates are orthonormal in $\mathcal{H}_d$, Euclidean parameter distance equals function-space distance. If the population best fit additionally satisfies $\|f_{d,k}^\circ - f_d^*\|_d^2 \leq C_{\text{app}} B_d(k)$, the squared triangle inequality gives equation D.1 directly.

If

$$p^G := \sup_d p_d^G < \infty,$$

then $p_d^G \leq p^G$ for every d , and therefore

$$\left\| \hat{\theta}_{n,d}^G - \theta_d^* \right\|_2^2 \leq C \frac{\sigma^2 p^G + \log(1/\delta)}{\alpha^2 n}.$$

F.3 PROOF OF THE TRANSFER BOUND

Fix a source dimension d . Suppose that for every $D \geq d$,

$$\Phi_{d \rightarrow D} \theta_d^* = \theta_D^*,$$

and that $\Phi_{d \rightarrow D}$ is distance preserving on the invariant model:

$$\|\Phi_{d \rightarrow D} \theta - \Phi_{d \rightarrow D} \theta'\|_2 = \|\theta - \theta'\|_2 \quad \forall \theta, \theta' \in \Theta_d^{G_d}.$$

Then, for every $D \geq d$,

$$\begin{aligned} \left\| \Phi_{d \rightarrow D} \hat{\theta}_{n,d}^G - \theta_D^* \right\|_2 &= \left\| \Phi_{d \rightarrow D} \hat{\theta}_{n,d}^G - \Phi_{d \rightarrow D} \theta_d^* \right\|_2 \\ &= \left\| \hat{\theta}_{n,d}^G - \theta_d^* \right\|_2. \end{aligned}$$

Taking the supremum over $D \geq d$, we get

$$\sup_{D \geq d} \left\| \Phi_{d \rightarrow D} \hat{\theta}_{n,d}^G - \theta_D^* \right\|_2^2 = \left\| \hat{\theta}_{n,d}^G - \theta_d^* \right\|_2^2.$$

Combining this identity with the estimation bound above yields

$$\sup_{D \geq d} \left\| \Phi_{d \rightarrow D} \hat{\theta}_{n,d}^G - \theta_D^* \right\|_2^2 \leq C \frac{\sigma^2 p_d^G + \log(1/\delta)}{\alpha^2 n}$$

with probability at least $1 - \delta$. If $p^G = \sup_d p_d^G < \infty$, the right-hand side is bounded by

$$C \frac{\sigma^2 p^G + \log(1/\delta)}{\alpha^2 n}.$$

This completes the auxiliary transfer calculation.

G EQUIVARIANT FIELD SPECIALIZATION FOR SETS

Score matching and denoising replace invariant scalar functions by permutation-equivariant vector fields, but the proof is otherwise unchanged. Let $\{\psi_{d,j}\}_{j \geq 1}$ be an orthonormal equivariant basis and write

$$s_d^* = \sum_{j \geq 1} \theta_{d,j}^* \psi_{d,j}, \quad \|\hat{s}_d - s_d^*\|_d^2 = \sum_{j \geq 1} (\hat{\theta}_{d,j} - \theta_{d,j}^*)^2.$$

If $\mathcal{I}_{d,k}$ contains the features of degree at most k , then its stable dimension and the omitted coefficient energy are

$$m_r(k) = |\mathcal{I}_{d,k}| = \exp\left(C_r k^{r/(r+1)} + o\left(k^{r/(r+1)}\right)\right), \quad B_d(k) = \sum_{j \notin \mathcal{I}_{d,k}} (\theta_{d,j}^*)^2.$$

Orthogonality and the auxiliary finite-dimensional bound in Appendix F give, with probability at least $1 - \delta$,

$$\|\hat{s}_{n,d}^{(k)} - s_d^*\|_d^2 \leq B_d(k) + C \frac{\sigma^2}{\alpha^2} \frac{m_r(k) + \log(1/\delta)}{n}. \quad (\text{G.1})$$

Thus the scalar proof applies verbatim. Uniform polynomial decay $B_d(k) \lesssim k^{-\tau}$ gives $(\log n)^{-\tau(r+1)/r}$. Under uniform exponential decay $B_d(k) \lesssim e^{-ak^\nu}$, with $\beta = r/(r+1)$,

$$\inf_k \left\{ B_d(k) + \frac{m_r(k)}{n} \right\} \asymp \begin{cases} n^{-1+o(1)}, & \nu > \beta, \\ n^{-a/(a+C_r)+o(1)}, & \nu = \beta, \\ \exp\left[-aC_r^{-\nu/\beta} (\log n)^{\nu/\beta} (1+o(1))\right], & 0 < \nu < \beta. \end{cases} \quad (\text{G.2})$$

Because the equivariant transfer operator is isometric, these bounds hold simultaneously at every target size $D \geq d$.

H EQUIVARIANT FIELD SPECIALIZATION FOR GRAPHS

Let $\mathcal{J}_{d,k}$ contain the degree-at-most- k equivariant graph features, and write $M_r(k) = |\mathcal{J}_{d,k}|$. For fixed channel dimension r ,

$$M_r(k) = \exp(k \log k + O_r(k \log \log k)), \quad B_d^{\text{graph}}(k) = \sum_{j \notin \mathcal{J}_{d,k}} (\theta_{d,j}^*)^2.$$

The same orthogonal decomposition and finite-dimensional estimate yield

$$\|\hat{s}_{n,d}^{(k)} - s_d^*\|_d^2 \leq B_d^{\text{graph}}(k) + C \frac{\sigma^2}{\alpha^2} \frac{M_r(k) + \log(1/\delta)}{n}. \quad (\text{H.1})$$

Consequently, polynomial decay $B_d^{\text{graph}}(k) \lesssim k^{-\tau}$ gives $(\log n / \log \log n)^{-\tau}$, while exponential decay $B_d^{\text{graph}}(k) \lesssim e^{-ak^\nu}$ gives

$$\inf_k \left\{ B_d^{\text{graph}}(k) + \frac{M_r(k)}{n} \right\} \asymp \begin{cases} n^{-1+o(1)}, & \nu > 1, \\ \exp\left[-a \left(\frac{\log n}{\log \log n}\right)^\nu (1+o(1))\right], & 0 < \nu \leq 1. \end{cases} \quad (\text{H.2})$$

The channel dimension affects only the lower-order term in $M_r(k)$, so it does not enter these first-order rates. Isometric equivariant transfer again makes every bound simultaneous over $D \geq d$.

I EXPERIMENTS

The experiments isolate two complementary parts of the theory rather than attempting a broad architectural benchmark. The first asks whether an invariant representation learned at one graph size retains its meaning at larger sizes. The second asks what remains when transfer is exact but the target must be approximated by a low-degree truncation. Together they separate representation-induced transfer error from approximation error due to the source tail.

I.1 SUBGRAPH COUNTING

We include a small synthetic experiment illustrating the role of invariant low-degree structure in transfer across graph sizes. The purpose of the experiment is not to benchmark a new architecture, but to visualize the basic mechanism behind our theory: a low-degree invariant statistic has a stable meaning across graph sizes, while a non-invariant coordinate polynomial does not transfer canonically.

We generate graphs from an Erdős–Rényi model with a graph-dependent edge probability. For each sample, we draw

$$p \sim \text{Unif}(0.15, 0.85), \quad A \sim G(d, p),$$

where $A \in \{0, 1\}^{d \times d}$ is the adjacency matrix of an undirected graph with no self-loops. The response is the population triangle density $y \approx p^3$. Thus, the target is a low-degree graph moment, but the learner observes only the sampled graph A , not the latent parameter p . We train all models only on graphs with $d = 10$ nodes and evaluate them on larger graphs with

$$D \in \{25, 30, 35, 40, 50, 60, 75, 100, 125, 150, 175, 200\}.$$

We compare two predictors. The first is an invariant polynomial model using normalized graph statistics

$$\phi_{\text{inv},d}(A) = (E_d(A), W_d(A), T_d(A)),$$

where $E_d(A)$ is the edge density, $W_d(A)$ is the normalized wedge density, and $T_d(A)$ is the normalized triangle density. A linear model is fit on these features at $d = 10$, and transfer to size D is performed by evaluating the same normalized invariant statistics on the larger graph:

$$\hat{f}_{\text{inv},D}(A) = \hat{\beta}_0 + \hat{\beta}_1 E_D(A) + \hat{\beta}_2 W_D(A) + \hat{\beta}_3 T_D(A).$$

This gives a direct size-transfer rule because the features are invariant graph moments whose interpretation is stable across graph sizes.

The second predictor is a non-invariant degree-three polynomial baseline. At the training size $d = 10$, we use the raw triangle indicators

$$A_{ij}A_{ik}A_{jk}, \quad 1 \leq i < j < k \leq 10,$$

as features and fit a linear model. Such a model assigns separate coefficients to labeled triangles and therefore has no canonical extension to graphs with more than ten nodes. To give this baseline a favorable transfer rule, we average it over random induced subgraphs: for a test graph on D nodes, we sample subsets $S_1, \dots, S_M \subseteq [D]$ of size ten and define

$$\hat{f}_{\text{raw},D}^{\text{avg}}(A) = \frac{1}{M} \sum_{m=1}^M \hat{f}_{\text{raw},10}(A_{S_m \times S_m}).$$

This is a Monte Carlo symmetrization of the raw non-invariant predictor. It deliberately gives the baseline an advantageous test-time correction: without averaging over induced subgraphs, its coordinates refer only to the ten training labels and do not define a predictor on a larger graph. The comparison therefore measures whether invariance is built into the learned representation or reconstructed afterwards by explicit averaging.

Figure 1(a) shows the test mean-squared error as the test graph size increases. The invariant polynomial model transfers directly and improves with graph size, since its normalized moments use all available nodes and become more accurate estimates of the underlying population quantities. The raw model also improves after subgraph averaging, but remains consistently worse despite this favorable correction. Thus, low polynomial degree alone is not the operative property: transfer comes from features whose invariant meaning is stable across sizes. The experiment also clarifies that larger test graphs need not be intrinsically harder; once the representation is size-consistent, additional nodes can provide more information about the same latent graph statistics.


I.2 EQUIVARIANT FIELD TRANSFER ON WEIGHTED GRAPHS

We next illustrate equivariant field estimation in a controlled setting where the target is known analytically. Continuous weighted graphs are represented by symmetric matrices

$$A \in \mathbb{R}^{d \times d},$$

with zero diagonal and independent standard-Gaussian upper-triangular entries. We regress onto the permutation-equivariant field

$$s_d^*(A) = -A - \sum_{q=2}^8 b_q q d^{-q} A^{q-1}.$$



(a) Invariant versus symmetrized raw features.

(b) Fast versus slow coefficient decay.

Figure 1: Two mechanisms governing transfer across graph sizes. **(a)** Transfer from 10-node Erdős–Rényi graphs: normalized invariant moments transfer directly, whereas labeled triangle features require explicit subgraph averaging. **(b)** Exact equivariant transfer from 10-node weighted graphs using size-normalized features: curves show mean relative MSE over ten repetitions, with 95% confidence intervals. Fast coefficient decay yields uniformly smaller truncation error.

The features $qd^{-q}A^{q-1}$ are gradients of normalized invariant spectral statistics $d^{-q}\text{tr}(A^q)$. Their meaning is shared across graph sizes, and the same coefficient b_q therefore defines the same rule at every size. We compare a fast-decay target,

$$b_q = 2 \exp(-0.14q^2),$$

with a slow-decay target,

$$b_q = 2q^{-1/2},$$

for which the omitted high-order components remain substantial.

For each regime, we fit the degree-three equivariant model

$$\hat{s}_d(A) = -A - \sum_{q=2}^4 \hat{b}_q qd^{-q}A^{q-1}$$

by least squares on the off-diagonal field entries of 200 independently sampled graphs with $d = 10$. Transfer to size D reuses the learned coefficients $\hat{b}_q$ and evaluates the corresponding normalized features $qD^{-q}A^{q-1}$. We test on 16 independent graphs at each

$$D \in \{10, 15, 20, 30, 40, 60, 80\}.$$

Because the common linear term $-A$ is represented exactly, we report error relative to the energy of the nonlinear target component:

$$\text{RelMSE}_D = \frac{\sum_A \sum_{i < j} (\hat{s}_D(A)_{ij} - s_D^*(A)_{ij})^2}{\sum_A \sum_{i < j} (s_D^*(A)_{ij} + A_{ij})^2}.$$

The full procedure is repeated for ten independent seeds, and the shaded bands in Figure 1(b) report 95% confidence intervals across these repetitions. No parameters are refit or calibrated at the target sizes: the only size-dependent operation is evaluating the analytically normalized features at (D).

Figure 1(b) shows that the same learned rule remains stable across all tested sizes. The absence of systematic deterioration with (D) is the empirical signature of exact equivariant transfer in this example. What remains is truncation error: fast decay makes the degree-three approximation highly accurate, whereas slow decay leaves a persistent gap roughly two to three orders of magnitude larger. This separates the two roles appearing in the theory—normalization controls size mismatch, while coefficient decay determines how much signal lies beyond the fitted degree. The experiment is illustrative rather than a test of the asymptotic minimax rates, which additionally depend on the full growth of invariant feature spaces.

J NOTATION

Symbol	Meaning
d, D	Training object size and target object size, with $D \geq d$.
n	Number of training samples observed at size d .
r	Element-feature dimension for sets or channel dimension for attributed graphs.
$\mathcal{X}_d$	Size- d input space: $\mathbb{R}^{r \times d}$ for sets or $\mathbb{R}^{r \times d \times d}$ for graphs.
Z_i	Unified observation: X_i for density estimation and (X_i, Y_i) for regression.
λ_d	Standard Gaussian reference measure on $\mathcal{X}_d$.
μ_d, ρ_d	Input distribution and joint distribution of (X, Y) , respectively.
f_d^*	Population target at size d : a density relative to λ_d or a regression function.
$\mathcal{H}_d$	$L^2(\lambda_d)$ for density estimation or $L^2(\mu_d)$ for regression.
$\ \cdot\ _d$	The corresponding norm on $\mathcal{H}_d$.
$\mathcal{F}_d, \mathcal{F}_d^{\text{inv}}$	Statistical function class and its permutation-invariant subclass.
$\ell_d, \hat{L}_{n,d}, L_d$	Pointwise loss, empirical risk, and population risk.
S_d	Symmetric group acting by relabeling set elements or graph nodes.
$\hat{f}_{n,d}^{\text{inv}}$	Invariant empirical risk minimizer learned at size d .
$T_{d \rightarrow D}$	Exact coherent isometric transfer operator from size d to size D .
k	Polynomial degree or interaction order.
$\mathcal{V}_{d,k}$	Invariant polynomials of degree at most k in $\mathcal{H}_d$.
$\mathcal{F}_{d,k}$	Order- k approximation class; it also enforces density admissibility when required.
$f_{d,k}^\circ, \hat{f}_{n,d,k}^{\text{inv}}$	Population and empirical risk minimizers over $\mathcal{F}_{d,k}$.
$B_d(k), B(k)$	Best order- k approximation error at size d and its supremum over sizes.
$q_X(k)$	Stable dimension $\sup_d \dim(\mathcal{V}_{d,k})$ for $X \in \{\text{set}, \text{graph}\}$.
$N_r(k), G_r(k)$	Stable homogeneous degree- k invariant-feature counts for sets and graphs.
τ	Polynomial source-decay exponent: $B(k) \lesssim k^{-\tau}$.
a, ν	Exponential source-decay constant and exponent: $B(k) \lesssim e^{-ak^\nu}$.
β	Set representation-growth exponent $r/(r+1)$.
C_r	Set growth constant $\frac{r+1}{r} (r\zeta(r+1))^{1/(r+1)}$.
δ	Failure probability in high-probability bounds.
$\mathfrak{F}_X(B)$	Size-consistent source class used in the minimax theorem.
$\mathfrak{R}_n(X, B)$	Worst-case minimax size-generalization risk.
$s_d^*, \theta_{d,j}^*$	Equivariant field target and its coefficients in the optional score or denoising specialization.
$\mathcal{I}_{d,k}, \mathcal{J}_{d,k}$	Degree-at-most- k equivariant feature indices for sets and graphs.
$m_r(k), M_r(k)$	Stable dimensions of the set and graph equivariant feature spaces.
α, σ	Uniform curvature constant and gradient-noise scale in the auxiliary estimation bound.
C, c, c_i	Positive constants independent of n, d, D ; their values may change between displays.