

WEATHERBRIDGE: FORECAST-AWARE MULTI-SOURCE INPUT ADAPTATION FOR PRETRAINED WEATHER FORECASTING MODELS

Mutian Xi^{1,2*}, Xinyu Su^{1,2*}, Xiaohui Zhong^{1,2,3†} & Hao Li^{1,2,3†}

¹Artificial Intelligence Innovation and Incubation Institute, Fudan University

²Shanghai Academy of AI for Science

³FuXi Intelligent Computing Technology Co., Ltd.

ABSTRACT

Data-driven weather forecasting models are commonly trained on ERA5, yet operational deployment must use timely initial states from physics-based numerical weather prediction (NWP) systems whose distributions differ substantially from ERA5. Direct initialization with these states can degrade autoregressive forecasts, motivating input adaptation that keeps the pretrained forecasting model unchanged. Here, we introduce WeatherBridge, a unified, source-conditioned adapter that maps heterogeneous operational NWP initial states to ERA5 reference states without updating the pretrained forecasting model. WeatherBridge predicts the structured residual corrections with a Swin-UNet and injects source embeddings across multiple spatial scales, allowing the model to capture shared atmospheric structure and source-specific discrepancies. Its training objective combines latitude-weighted ERA5 reconstruction with One-step Rollout Alignment (ORA), which backpropagates one-step forecast error through a frozen FuXi model to optimize the adapted states for both initial state fidelity and downstream forecast compatibility. We evaluate WeatherBridge using daily GFS, HRES, and CMA initializations from 1 January to 22 November 2025 across six variables and lead times up to 240 hours. Relative to raw source initialization of FuXi, WeatherBridge reduces the mean RMSE of FuXi over the 6- 240-hour rollout for 17 of 18 source-variable combinations, with a median reduction of 10.2% and a maximum reduction of 46.3%. The adapter trained with frozen FuXi also improves FengWu forecasts on GFS inputs without retraining either the adapter or FengWu. These results demonstrate that forecast-aware input adaptation can improve forecasts from heterogeneous operational NWP inputs and transfer across pretrained ERA5-based forecasting models.

1 INTRODUCTION

Data-driven weather forecasting models have emerged as promising alternatives to traditional numerical weather prediction (NWP) models for global medium-range weather forecasting. State-of-the-art models learn atmospheric evolution from historical data and can match or outperform conventional NWP models at substantially lower inference cost (Lam et al., 2023; Bi et al., 2023; Chen et al., 2025; 2023; Lam et al., 2023; Bodnar et al., 2025). Most of these models are trained on ERA5, whose data quality, global coverage, temporal consistency, public availability, and comprehensive set of multilevel variables have made it a standard benchmark for data-driven weather forecasting (Hersbach et al., 2020). However, operational forecasting requires timely initial conditions from NWP systems such as GFS, HRES, and CMA because ERA5 is available only after a delay of approximately five days. Differences in data assimilation and numerical modeling cause these operational fields to exhibit statistical and structural characteristics that differ substantially from those of ERA5. Consequently, directly initializing an ERA5-trained model with operational NWP fields can significantly degrade forecast performance.

*These authors contributed equally to this work.

†Corresponding author. zhong_xiaohui@fudan.edu.cn, lihao_lh@fudan.edu.cn




Figure 1: Effect of operational-source initialization on a frozen ERA5-trained forecaster. (a) Schematic illustration of source-dependent input distribution differences. (b) Initialization of the same frozen forecaster with ERA5 or raw operational fields, without adaptation. (c) T2M RMSE of FuXi initialized with ERA5, GFS, HRES, and CMA, with all forecasts verified against ERA5. The forecast errors vary across initialization sources.

This input distribution shift is especially consequential for autoregressive forecasting. Models such as FuXi, GraphCast, and Pangu-Weather implicitly rely on the variable definitions, normalization statistics, spatial structures, and multivariate relationships in ERA5. Initial fields from GFS, HRES, and CMA exhibit source-specific discrepancies in these characteristics. When such fields are fed directly to a frozen ERA5-trained model, the model operates outside its training distribution, and differences in the initial state can propagate and amplify through the autoregressive rollout. Figure 1 illustrates this effect for FuXi. Initialization with raw GFS or CMA yields higher RMSE than ERA5 initialization at all lead times on variable 2 meter temperature (t2m). HRES initialization begins closer to the ERA5 reference, and the performance gap generally narrows at longer lead times as forecast errors saturate. Therefore, the extent of the degradation depends strongly on the discrepancy between each operational NWP source and ERA5.

Fine-tuning is one approach to adapting pretrained forecasting models to different initialization data. For example, FuXi-GRAPES fine-tunes FuXi on CMA reanalysis data (Zhang et al., 2025), while operational GraphCast is fine-tuned on ECMWF HRES analysis (Google DeepMind, 2024). Changes to operational systems can also prompt further training, as illustrated by AIFS v2 following the IFS Cycle 50r1 upgrade (ECMWF, 2026b;a). We explore a complementary approach that trains a shared input adapter while keeping the pretrained forecasting model unchanged.

We introduce WeatherBridge, a unified, source-conditioned adapter that maps heterogeneous operational NWP fields to ERA5 reference states while leaving the pretrained weather forecasting models unchanged. WeatherBridge represents the NWP-to-ERA5 mapping as a structured residual correction learned by a shared Swin-UNet. To account for systematic differences among GFS, HRES, and CMA, WeatherBridge injects a learnable source embedding into the network at multiple spatial scales. This hierarchical conditioning enables a single model to capture shared atmospheric structure while retaining NWP-specific corrections.

Crucially, matching ERA5 at the initial time does not by itself guarantee that an adapted state will support accurate autoregressive forecasts. Therefore, WeatherBridge combines latitude-weighted ERA5 reconstruction with One-step Rollout Alignment (ORA). ORA propagates one-step forecast error through a frozen forecasting model, encouraging the adapter to produce states that are compatible with the model’s learned dynamics and close to the corresponding ERA5 fields. In this study, we train WeatherBridge using the frozen FuXi model and evaluate the resulting adapter on daily GFS, HRES, and CMA initializations over lead times of up to 240 hours. Relative to direct initialization with raw source fields, WeatherBridge reduces FuXi’s mean rollout RMSE for 17 of the 18 source-variable combinations, with reductions of up to 46.3%. When applied unchanged to the independently trained FengWu model, WeatherBridge also reduces RMSE and increases ACC for all six evaluated GFS variables at both 6 and 240 hours.

Our contributions are as follows:

- We propose WeatherBridge, a unified, source-conditioned adapter that maps heterogeneous operational NWP initial fields to ERA5 reference states, allowing a frozen ERA5-trained forecasting model to use non-ERA5 inputs without fine-tuning.
- We introduce a hierarchical source conditioning mechanism that injects learned source embeddings at multiple spatial scales, allowing a single shared adapter to account for source-dependent discrepancies when mapping operational NWP inputs to ERA5 reference states.
- We introduce a dual-target training objective that combines latitude-weighted ERA5 reconstruction with One-step Rollout Alignment (ORA) through the frozen forecasting model, jointly promoting initial state fidelity and downstream forecast performance.
- We evaluate WeatherBridge on three operational NWP sources (GFS, HRES, CMA), six variables, and forecast lead times up to 240 h. The adapter improves FuXi forecasts and transfers unchanged to FengWu (another ERA5-trained forecasting model).

2 RELATED WORK

ERA5-Trained AI-based Weather Forecasting Models. Recent AI-based systems, such as GraphCast, Pangu-Weather, FuXi, FourCastNet, FengWu, GenCast, and AIFS, have shown competitive medium-range forecast skill by learning global atmospheric evolution from reanalysis or operational analysis data (Lam et al., 2023; Bi et al., 2023; Chen et al., 2023; Pathak et al., 2022; Chen et al., 2025; Price et al., 2025; Lang et al., 2024). Despite their architectural differences, these models generally perform most reliably when their initial states follow the distribution encountered during training. Therefore, operational deployment with a different NWP source may require source-specific adaptation. WeatherBridge addresses the distribution mismatch at the input interface by mapping heterogeneous operational initial states to the ERA5 reference domain while leaving the forecasting model unchanged.

Bias Correction and Statistical Post-Processing. An alternative to model fine-tuning is to correct systematic errors in NWP or AI weather forecasts using bias correction, model output statistics (MOS), quantile mapping, and AI-based post-processing. These methods typically target particular variables, locations, or products (Vannitsem et al., 2021; Huang et al., 2024). WeatherBridge transforms the initial state supplied to a pretrained forecasting model, with the objective of improving both its agreement with ERA5 and the resulting autoregressive forecasts. To this end, its training objective propagates forecast error through the frozen FuXi model, explicitly accounting for how the adapted state affects downstream forecast performance.

Earth-System Foundation Models and Heterogeneous Data. Earth-system foundation models such as Aurora and ClimaX accommodate multiple data sources, resolutions, variable sets, and prediction tasks through large-scale pretraining and task-specific adaptation (Bodnar et al., 2025; Nguyen et al., 2023). This strategy seeks to make the model broadly transferable by exposing it to heterogeneous data during training. WeatherBridge pursues a complementary pathway. Rather than requiring a single forecasting model to accommodate multiple NWP data distributions, it preserves the ERA5-trained backbone and learns a source-conditioned mapping from the operational NWP initial states to ERA5 reference state. The distinction is between building heterogeneity into the forecasting model and resolving heterogeneity before the model is used.

Model Reprogramming and Input-Side Adaptation. WeatherBridge is also related to model reprogramming, which redirects a frozen or lightly modified pretrained model to new tasks or domains through input transformations, prompts, masks, or lightweight modules (Ye et al., 2025; Cai et al., 2024). Both approaches keep the major pretrained model unchanged and adapt its interface to a new input distribution. However, WeatherBridge operates on physically meaningful, multivariate, spatio-temporally continuous meteorological states rather than image patches or tokens. In addition, WeatherBridge is trained not only to reconstruct a reference state but also to improve the subsequent autoregressive forecast.

3 METHODOLOGY

We present WeatherBridge, a plug-and-play adapter that transforms heterogeneous operational NWP initial states into ERA5-aligned states for a frozen pretrained weather forecasting model. In this work, we instantiate the forecasting backbone with FuXi (Chen et al., 2023) while keeping its architecture and parameters unchanged throughout adapter training. Figure 2 illustrates the overall framework. We first formalize the input adaptation problem in Section 3.1. We then describe the source-conditioned adapter architecture in Section 3.2, which builds on a residual Swin-UNet and injects source identity hierarchically at multiple feature resolutions. Finally, we introduce the forecast-aware training strategy in Section 3.3, which combines latitude-weighted ERA5 reconstruction with One-step Rollout Alignment (ORA) while keeping weather forecasting backbone frozen.

3.1 PROBLEM FORMULATION

Let $\mathcal{S} = \{\text{GFS}, \text{HRES}, \text{CMA}\}$ denote the set of operational NWP sources. We index atmospheric states at 6-hour intervals, such that $t - 1$, t , and $t + 1$ denote three consecutive valid times. For each source $s \in \mathcal{S}$, let x_t^s and x_t^E denote the source and ERA5 states, respectively, at the same valid time t . Both are represented as tensors in $\mathbb{R}^{C \times H \times W}$, where $C = 70$ is the number of atmospheric channels and $(H, W) = (721, 1440)$ correspond to the latitude and longitude dimensions of the global 0.25° .

Given a single source state, its source identity, and the associated temporal information, WeatherBridge learns the frame-wise mapping

$$\hat{x}_t^E = F_\theta^{\text{ada}}(x_t^s, s, t), \quad (1)$$

where $\hat{x}_t^E$ is the reference state produced from the source state at time t . The adapter always operates on one atmospheric state at a time.

Let F_ϕ denote a forecasting model pretrained on ERA5, whose parameters ϕ remain fixed. Given two consecutive source states, the goal is to learn a frame-wise adapter F_θ^{ada} such that the states obtained by independently adapting the two frames enable F_ϕ to accurately predict the subsequent atmospheric evolution. Formally, the desired property is

$$\tilde{x}_{t+k}^E = F_\phi^{[k]} \left(\left[F_\theta^{\text{ada}}(x_{t-1}^s, s, t-1), F_\theta^{\text{ada}}(x_t^s, s, t) \right] \right) \approx x_{t+k}^E, \quad k = 1, \dots, K, \quad (2)$$

where $[\cdot, \cdot]$ denotes temporal concatenation, and $F_\phi^{[k]}$ denotes the atmospheric state predicted for valid time $t + k$ through a k -step autoregressive rollout of the frozen forecasting model.

3.2 SOURCE-CONDITIONED ADAPTER ARCHITECTURE

Figure 2 provides an overview of WeatherBridge and its integration with the frozen pretrained forecasting model. WeatherBridge employs a residual Swin-UNet to transform each source state into the reference state. A hierarchical conditioning mechanism injects source identity at multiple feature resolutions, enabling the shared adapter to learn source-specific corrections across spatial scales.

Residual Swin-UNet Backbone. Our adapter employs a residual Swin-UNet backbone to transform each source state into the reference state. A convolutional patch embedding first projects the atmospheric variables into a latent feature space. The encoder stacks convolutional residual blocks with strided downsampling to progressively aggregate spatial information, while a window-based self-attention block at the bottleneck models broader spatial dependencies at a reduced resolution. The decoder uses interpolation-based upsampling and convolutional residual blocks to reconstruct the atmospheric state. At each decoder stage, the upsampled representation is fused with the corresponding encoder feature through a skip connection, retaining spatial information that may be weakened during downsampling. A patch-recovery head then maps the decoded representation back to the original variable space and spatial grid. Rather than predicting the complete target state directly, the network estimates a residual correction to the normalized source field:

$$\hat{x}_t^E = F_\theta^{\text{ada}}(x_t^s, s, t) = x_t^s + \Delta_\theta(x_t^s, s, t), \quad (3)$$

where Δ_θ denotes the correction predicted by the encoder-decoder backbone. Both x_t^s and $\Delta_\theta(x_t^s, s, t)$ are represented in the same normalized variable space, and conversion back to physical units is performed outside the adapter when required. This residual parameterization provides a direct identity path from the source state to the adapted state and represents the source-to-ERA5 transformation as a correction to the original field.




Figure 2: Overview of WeatherBridge. (a) *Forecast-aware training*. Given a source state x_t^s , WeatherBridge maps it to a reference-domain state $\hat{x}_t^E$. The adapted state and the preceding ERA5 state x_{t-1}^E are fed into the frozen weather forecasting model to predict $\hat{x}_{t+1}^E$. The adapter is optimized using the reconstruction loss $\mathcal{L}_{\text{rec}}$ at time t and the one-step rollout loss $\mathcal{L}_{\text{ORA}}$ at time $t+1$, while the forecasting model remains frozen. (b) *Inference stage*. Two consecutive source states, x_{t-1}^s and x_t^s , are mapped by WeatherBridge with shared parameters. The resulting state pair is then passed to the frozen weather forecasting model to predict $\hat{x}_{t+1}^E$. (c) *WeatherBridge adapter*. The source identity and valid time are encoded as embeddings and hierarchically injected into multiple downsampling stages of a residual Swin-UNet. The network predicts a source-aware and time-aware correction $\Delta_\theta(x_t^s, s, t)$, yielding $\hat{x}_t^E = x_t^s + \Delta_\theta(x_t^s, s, t)$.

Hierarchical Conditioning. WeatherBridge shares the same encoder–decoder parameters across all NWP sources while conditioning the network on the identity of each source. Specifically, the categorical source identifier $s \in \mathcal{S}$ is used to retrieve a source embedding from a learnable lookup table, i.e., $e_s = E_{\text{src}}[s]$. Here, $E_{\text{src}} \in \mathbb{R}^{|\mathcal{S}| \times d_s}$, and d_s denotes the embedding dimension. This embedding table is optimized jointly with the adapter, enabling a single shared backbone to retain source-specific information.

In addition to the source identity, we derive a fixed sinusoidal temporal embedding e_t from the seasonal phase of the valid time, computed using the day of year and hour of day. This embedding provides temporal metadata associated with the current atmospheric state and does not introduce an additional input frame.

After being projected to the input channel dimension, the source and temporal embeddings are broadcast over the spatial grid and added to the source field before patch embedding. To preserve the conditioning information throughout the encoder hierarchy, both embeddings are projected and injected at every downsampling level. Given a downsampling feature h_l at level l , the conditioned representation is computed as

$$\bar{h}_l = h_l + \mathcal{P}_l^s(e_s) + \mathcal{P}_l^t(e_t), \quad (4)$$

where $\mathcal{P}_l^s$ and $\mathcal{P}_l^t$ denote stage-specific projections that match the channel dimension and spatial resolution of h_l . Conditioning is applied throughout the encoder but not in the decoder, allowing the shared backbone to model source-dependent corrections at multiple spatial scales.

3.3 FORECAST-AWARE TRAINING STRATEGY

We train WeatherBridge to map heterogeneous NWP source fields to adapted initial fields that support downstream forecasting. Toward this goal, we use the standard reference-state reconstruction loss $\mathcal{L}_{\text{rec}}$ as the base training objective, encouraging the adapted initial field to approach its paired reference state. However, reconstruction alone does not explicitly account for how the adapted field affects the downstream forecasting model. We therefore augment the base objective with our proposed One-step Rollout Alignment (ORA) loss $\mathcal{L}_{\text{ORA}}$, which propagates the one-step forecast error through the frozen forecasting model. The resulting training objective is

$$\mathcal{L}_{\text{total}} = \mathcal{L}_{\text{rec}} + \lambda_f \mathcal{L}_{\text{ORA}}, \quad (5)$$

where $\lambda_f > 0$ controls the contribution of the forecast-aware ORA term.

Specifically, the reconstruction term is defined as a latitude-weighted ℓ_1 loss between the adapted initial field and its paired ERA5 reference state:

$$\mathcal{L}_{\text{rec}} = \frac{1}{C} \sum_{c=1}^C \frac{\sum_i w_i |\hat{x}_{t,i}^{(c)} - x_{t,i}^{E,(c)}|}{\sum_i w_i}, \quad w_i = \cos(\phi_i), \quad (6)$$

where C is the number of atmospheric channels, i indexes spatial grid points, and ϕ_i denotes the latitude of grid point i . The latitude weights correct for unequal grid-cell areas, with both $\hat{x}_t$ and x_t^E represented in normalized space.

To introduce forecast-aware supervision, ORA evaluates the adapted initial field through the frozen forecasting model. Given the adapter output $\hat{x}_t^E$, we combine it with the preceding ERA5 reference state to construct the two-frame input $[x_{t-1}^E, \hat{x}_t^E]$. The frozen model then produces the one-step forecast $\tilde{x}_{t+1}$. Using the ERA5 reference state as the preceding frame isolates the effect of the current adapted field without introducing an additional error from the previous input frame. The forecast error is back-propagated through F_ϕ to update the adapter parameters only.

For each guided channel, we first compute a latitude-weighted RMSE over the spatial grid and then normalize it by the channel-specific reference RMSE of FuXi under native ERA5 inputs. The resulting normalized errors are averaged equally across the guided channels:

$$\mathcal{L}_{\text{ORA}} = \frac{1}{|\mathcal{C}_{\text{ORA}}|} \sum_{c \in \mathcal{C}_{\text{ORA}}} \frac{\text{RMSE}_w^c(\tilde{x}_{t+1}^E, x_{t+1}^E)}{\text{RMSE}_{\text{ref}}^c}. \quad (7)$$

where RMSE_w^c is the latitude-weighted RMSE for channel c , and $\text{RMSE}_{\text{ref}}^c$ is obtained by forecasting from the native ERA5 input $[x_{t-1}^E, x_t^E]$. We apply ORA to

$$\mathcal{C}_{\text{ORA}} = \{\text{z500, z850, u200, t2m, u10m, v10m, msl, tp, t300}\}. \quad (8)$$

which spans upper-, middle-, lower-, and near-surface atmospheric conditions together with precipitation. The corresponding reference errors are reported in Table 4 in the Appendix.

At inference, the reference states are neither available nor required. WeatherBridge is applied independently to two consecutive states x_{t-1}^s and x_t^s from the same operational source, producing $\hat{x}_{t-1}^E$ and $\hat{x}_t^E$, respectively. The two adapted initial fields are then used as the FuXi input $[\hat{x}_{t-1}^E, \hat{x}_t^E]$. Thus, unlike ORA training, where the preceding frame is provided by the ERA5 reference sequence, both frames used for inference are derived exclusively from operational source data.

4 EXPERIMENTS

4.1 EXPERIMENTAL SETUP

Data. We use ERA5 as the target domain and GFS, HRES, and CMA as heterogeneous source domains. Each source field is paired with the ERA5 field at the same valid time. All datasets are represented as FuXi-compatible 70-channel global fields Chen et al. (2023) on a 0.25° grid with 721×1440 points, including 5 variables at 13 pressure levels and 5 surface variables. All input fields are normalized using the ERA5 statistics μ_{ERA5} and σ_{ERA5} . For all datasets, samples are initialized once per day, with consecutive forecast states spaced at 6-hour intervals. We use data from 2022-01-01 to 2024-06-30 for training, from 2024-07-01 to 2024-12-31 for validation, and from 2025-01-01 to 2025-11-22 for testing.

Implementation details. WeatherBridge contains 138.44M trainable parameters and uses a dropout rate of 0.1. With the FuXi backbone frozen, we train WeatherBridge in PyTorch on four A100 GPUs using FSDP, mixed precision, and AdamW with $\beta_1 = 0.9$, $\beta_2 = 0.999$, and a weight decay of 2×10^{-5} . We use a global batch size of 32, with 8 samples per GPU, clip the gradient norm at 5.0, linearly warm up the learning rate to 1×10^{-4} over the first 5 epochs, and then apply cosine decay to 1×10^{-7} over the remaining 85 epochs.




Figure 3: Forecasting performance on FuXi over the 240-hour rollout for the six evaluation variables. Higher ACC and lower RMSE indicate better forecasting performance. Solid curves denote the WeatherBridge-aligned initialization, while dashed curves of the same color denote the raw-source initialization for GFS (orange), HRES (green), and CMA (purple). The gray dashed curve denotes the ERA5 reference initialization.

Evaluation. Each test initialization is rolled out for 40 six-hour steps, corresponding to a forecast horizon of 240 hours. We compare three initialization paths: ERA5 $\rightarrow$ FuXi as the reference, raw source $\rightarrow$ FuXi as the direct-inference baseline, and raw source $\rightarrow$ WeatherBridge $\rightarrow$ FuXi as the proposed approach. We do not retrain or fine-tune FuXi separately for individual sources, as full source-specific retraining would require updating all 1,556.72M parameters of the backbone for each source and is outside the scope of this work. At inference, the proposed path uses only operational source fields. Following FuXi (Chen et al., 2023), all three initialization paths are evaluated against ERA5. We report results for $z500$, $t2m$, $t850$, $ws10$, $ws850$, and msl , where $ws10$ and $ws850$ are computed as the magnitudes of the horizontal wind vectors from the corresponding U and V components at 10 m and 850 hPa, respectively. We evaluate WeatherBridge at both the initialization and rollout forecasting levels. PSNR and SSIM provide supplementary measures of initial-state agreement with ERA5. Our primary evaluation uses latitude-weighted RMSE and ACC over the 240-hour rollout to assess downstream forecast skill. The definitions of all metrics are provided in Appendix A.3. We additionally report the ORA loss as a forecast-aware diagnostic for every configuration in the Appendix, and Table 4 lists the per-channel reference errors used to normalize it.

4.2 WEATHER FORECASTING PERFORMANCE

Figure 3 compares the ACC and RMSE trajectories produced by the three initialization paths over the 240-hour rollout. The benefit of WeatherBridge is most pronounced for GFS. Directly initializing FuXi with raw GFS fields leads to substantially lower ACC and higher RMSE across all six variables, with the gap generally increasing over the forecast horizon. After alignment, the GFS curves move consistently toward the ERA5 reference for both metrics, and the improvement persists throughout the rollout. CMA exhibits a similar but smaller improvement, with the clearest gains for T2M and MSL. In contrast, the raw and aligned HRES curves remain close to each other and to the ERA5 reference, indicating limited room for further improvement.

The mean rollout RMSE in Table 1 quantifies this source-dependent pattern. WeatherBridge reduces the mean RMSE for 17 of the 18 source-variable pairs. For GFS, the reductions range from 30.6% to 46.3% across the six variables. CMA also improves consistently, with reductions ranging from 4.3% to 26.4%, including the largest gains of 26.4% for T2M and 20.1% for MSL. The changes for HRES are considerably smaller: five variables show RMSE reductions of at most 1.04%, while Z500 exhibits a slight increase of 0.05%. Overall, the results show that WeatherBridge provides the

Table 1: Per-source and per-variable results on FuXi. PSNR and SSIM assess initial-state alignment with ERA5, while mean RMSE is averaged over the 40-step rollout up to 240 h. “↑” and “↓” indicate that higher and lower values are better, respectively. Δ denotes the relative improvement of the WeatherBridge-aligned result over the raw-source result; negative values indicate degradation.

Source	Var.	PSNR ↑			SSIM ↑			Mean RMSE ↓		
		raw	WeatherBridge	Δ	raw	WeatherBridge	Δ	raw	WeatherBridge	Δ
GFS	Z500	44.78	46.68	4.24%	0.9999	0.9999	0.00%	719.36	386.46	46.28%
	T2M	31.45	34.27	8.97%	0.9943	0.9971	0.28%	3.31	1.78	46.22%
	T850	29.61	33.68	13.75%	0.9922	0.9967	0.45%	3.60	2.11	41.39%
	WS10	21.89	23.77	8.59%	0.9409	0.9609	2.13%	3.09	2.08	32.69%
	WS850	23.50	24.18	2.89%	0.9529	0.9596	0.70%	5.06	3.51	30.63%
	MSL	23.95	35.34	47.56%	0.9442	0.9963	5.52%	642.47	372.90	41.96%
HRES	Z500	50.03	50.02	-0.02%	1.0000	1.0000	0.00%	338.17	338.33	-0.05%
	T2M	36.41	36.99	1.59%	0.9982	0.9985	0.03%	1.64	1.63	0.61%
	T850	36.17	36.81	1.77%	0.9981	0.9984	0.03%	1.92	1.90	1.04%
	WS10	24.94	25.43	1.96%	0.9711	0.9736	0.26%	1.93	1.92	0.52%
	WS850	25.72	25.98	1.01%	0.9715	0.9734	0.20%	3.23	3.23	0.00%
	MSL	39.09	39.60	1.30%	0.9985	0.9987	0.02%	330.32	329.38	0.28%
CMA	Z500	45.09	44.53	-1.24%	0.9998	0.9998	0.00%	446.81	403.29	9.74%
	T2M	30.87	33.61	8.88%	0.9936	0.9966	0.30%	2.50	1.84	26.40%
	T850	31.13	33.19	6.62%	0.9943	0.9965	0.22%	2.46	2.21	10.16%
	WS10	20.33	23.13	13.77%	0.9278	0.9544	2.87%	2.26	2.13	5.75%
	WS850	23.19	23.54	1.51%	0.9477	0.9528	0.54%	3.76	3.60	4.26%
	MSL	29.29	35.09	19.80%	0.9855	0.9963	1.10%	486.70	388.67	20.14%

Table 2: FengWu results on GFS at 240 h with the FuXi-trained adapter, neither retrained. ↑/↓: higher/lower is better. Δ : relative gain over raw-source initialization.

Var.	RMSE ↓			ACC ↑		
	Raw	Adapted	Δ (%)	Raw	Adapted	Δ (%)
Z500	1023.32	952.65	6.91	0.0732	0.1748	138.68
T2M	3.93	3.23	17.77	0.3584	0.4365	21.78
T850	4.81	4.08	15.25	0.0783	0.1717	119.27
WS10	3.88	3.86	0.61	0.0479	0.0907	89.50
WS850	6.29	5.91	6.08	0.0322	0.0873	171.17
MSL	867.90	727.29	16.20	0.0952	0.2912	205.81

greatest forecasting benefit when direct initialization causes a substantial loss of forecast skill, while largely preserving the performance of sources that are already compatible with the ERA5-trained backbone.

Transfer to FengWu. The adapter trained with frozen FuXi is applied directly to FengWu without retraining either the adapter or FengWu. On GFS inputs, adaptation reduces the 240-hour RMSE by 0.61–17.77% and increases ACC for all six evaluated variables (Table 2). Results at 6 h are provided in Appendix B.7.

4.3 INITIAL-STATE ALIGNMENT

Figure 4 shows the spatial distribution of the initial-state error reduction achieved by WeatherBridge relative to the raw-source fields, averaged over all 00Z initializations in the test period. Positive values indicate regions where the aligned field is closer to ERA5, while negative values indicate local degradation. For GFS, substantial error reductions appear across multiple variables, particularly for MSL and T2M. The Z500 map contains both pronounced improvements and degradations, showing that its aggregate improvement is not spatially uniform. CMA exhibits clear regional gains for T2M and WS10, whereas its Z500 error increases over broad areas. The HRES maps remain largely close to zero, apart from localized changes in T2M and Z500, consistent with the relatively small initial discrepancy between HRES and ERA5.




Figure 4: Reduction of the initial-field error against ERA5 obtained by the WeatherBridge-aligned initialization relative to the raw-source initialization, averaged over the 00z initializations of the test period. Rows are sources and columns are variables; red marks grid cells where the aligned initial field is closer to ERA5 than the raw-source field, blue marks cells where it is farther. The map compares initial fields rather than forecasts.

Table 1 summarizes the global alignment results. PSNR improves for all GFS variables, ranging from 2.89% to 47.56%, and for five CMA variables, ranging from 1.51% to 19.80%. The only notable decrease is CMA Z500 at 1.24%, while all HRES changes remain within 1.96%. SSIM improves in 15 of the 18 cases and remains unchanged in the other three. Together with the spatial maps, these results show improved initial-state agreement with ERA5 for GFS and CMA, limited changes for HRES, and occasional localized degradation.

4.4 ABLATION STUDY

We examine the effect of ORA in Figure 5. Compared with reconstruction-only training, ORA provides the largest RMSE reductions for MSL, followed by clear gains for T2M and Z500. Gains are smaller for T850 and WS850, with slight degradation at the first step, and generally diminish at longer lead times, consistent with the one-step ORA objective. Source-embedding ablations are provided in Appendix B.2, showing improved 6-hour RMSE for 13 of 18 source-variable combinations under balanced training and 17 under source imbalance.




Figure 5: NRMSE of GFS-only w/o ORA vs. GFS-only; negative (blue) means ORA helps.

Additional results. Further ablations, sensitivity tests, and analyses of significance, scaling, and efficiency appear in Appendix B.

5 CONCLUSION

We introduced WeatherBridge, a unified, source-conditioned adapter that maps heterogeneous operational NWP initial fields to ERA5 reference states while leaving the pretrained forecasting model unchanged. WeatherBridge represents the NWP-to-ERA5 mapping as a structured residual correction learned by a shared Swin-UNet, and injects a learnable source embedding at multiple spatial scales so that one adapter captures both shared atmospheric structure and source-specific discrepancies. Its training objective couples latitude-weighted ERA5 reconstruction with One-step Rollout Alignment, which propagates the forecast error of the frozen backbone back into the adapter, so that the adapted states are compatible with the dynamics learned by the backbone rather than merely close to ERA5, so the improvement is not confined to the single training step but holds over the

whole 6 h to 240 h rollout. However, ORA loss that applied to our adapter is one-step: the adapted state is optimized through a single forecast step, whereas forecast skill is ultimately judged over a long auto-regressive rollout. In future work, we plan to extend the alignment object to a multi-step rollout so that adapter is optimized against the error accumulation through long-lead forecasts.

AI USE STATEMENT

We used generative AI tools for language editing, LaTeX assistance, table formatting, and literature discovery for related work. We did not use generative AI tools to generate or alter data, run experiments, replace author decisions about the method, or make unsupported scientific claims. All AI-assisted text and code were reviewed by the authors. The authors designed the method and experiments, executed and tested the code, checked all numerical results against the recorded outputs, and manually verified the relevant literature. We take responsibility for the final content of this work, including any text, claims, code, or other artifacts produced with the aid of generative AI.

ETHICS STATEMENT

This work uses established meteorological reanalysis and numerical weather prediction datasets, including ERA5 and operational forecast sources, together with released pretrained weather forecasting models. It involves no human-subject data collection, personally identifiable information, or newly collected sensitive data. The datasets describe atmospheric and environmental conditions rather than individual-level information. Nevertheless, differences in spatial coverage, data quality, physical parameterizations, and source-model characteristics may introduce systematic biases into the learned alignment and downstream forecasts. We therefore recommend validating the proposed method on the target region, variables, and operational setting before deployment, particularly where forecasts may inform safety-critical decisions. We report the data sources, model configurations, and evaluation protocols used in this work to support reproducibility and responsible use.

REPRODUCIBILITY STATEMENT

The paper and appendix provide the complete method specification, the dataset and split descriptions, the preprocessing and normalization choices, the model and optimization settings, and the evaluation protocol, including the initialization paths, the forecast horizon, the verification variables and the metric definitions. All reported results follow a single deterministic evaluation protocol applied to the trained adapter weights, and the significance of the main improvements is established by a paired resampling test over the evaluation initializations. The pretrained forecasting models are used exactly as released and are never updated, both in the main experiments and in the cross-backbone transfer test, so no reported result requires retraining a forecasting model. Upon acceptance, we will release the complete source code, configuration files, preprocessing and evaluation scripts, the trained adapter for every configuration reported, and the run metadata needed to reproduce the experiments; the release will cover the multi-source adapter as well as the ablation, sensitivity and cross-backbone transfer settings.

REFERENCES

- Kaifeng Bi, Lingxi Xie, Hengheng Zhang, Xin Chen, Xiaotao Gu, and Qi Tian. Accurate medium-range global weather forecasting with 3d neural networks. *Nature*, 619:533–538, 2023.
- Cristian Bodnar, Wessel P. Bruinsma, Ana Lucic, et al. A foundation model for the earth system. *Nature*, 641:1180–1187, 2025. doi: 10.1038/s41586-025-09005-y.
- Chengyi Cai, Zesheng Ye, Lei Feng, Jianzhong Qi, and Feng Liu. Sample-specific masks for visual reprogramming-based prompting. In *Proceedings of the 41st International Conference on Machine Learning (ICML)*, pp. 5383–5408, 2024.
- K. Chen, T. Han, F. Ling, et al. The operational medium-range deterministic weather forecasting can be extended beyond a 10-day lead time. *Communications Earth & Environment*, 6:518, 2025. doi: 10.1038/s43247-025-02502-y.

- Lei Chen, Xiaohui Zhong, Feng Zhang, et al. Fuxi: A cascade machine learning forecasting system for 15-day global weather forecast. *npj Climate and Atmospheric Science*, 6:190, 2023.
- ECMWF. Adapting the aifs for 50r1 / implementation of aifs single v2, 2026a. URL <https://confluence.ecmwf.int/spaces/FCST/pages/620418870/Implementation+of+AIFS+Single+v2>. ECMWF Forecast User documentation; AIFS blog "Adapting the AIFS for 50r1": <https://www.ecmwf.int/en/about/media-centre/aifs-blog/2026/adapting-aifs-50r1>.
- ECMWF. Significant update to ecmwf's key forecasting systems ifs and aifs goes live (ifs cycle 50r1), 2026b. URL <https://confluence.ecmwf.int/spaces/FCST/pages/567162191/Implementation+of+IFS+Cycle+50r1>. ECMWF Forecast User documentation; news item (12 May 2026): <https://www.ecmwf.int/en/about/media-centre/news/2026/ifs-cycle-50r1-aifsv2-live>.
- Google DeepMind. Graphcast operational checkpoints. NVIDIA Earth2Studio documentation, 2024. URL <https://nvidia.github.io/earth2studio/main/modules/generated/models/px/GraphCastOperational/>. Operational checkpoint: pre-trained on ERA5 (1979–2017) and fine-tuned on ECMWF HRES analysis (2016–2021), initialized directly from HRES data.
- Hans Hersbach, Bill Bell, Paul Berrisford, et al. The era5 global reanalysis. *Quarterly Journal of the Royal Meteorological Society*, 146(730):1999–2049, 2020.
- Bohan Huang, Zhu Liu, Qingyun Duan, Adnan Rajib, and Jina Yin. Unsupervised deep learning bias correction of cmip6 global ensemble precipitation predictions with cycle generative adversarial network. *Environmental Research Letters*, 19:094003, 2024. doi: 10.1088/1748-9326/ad66e6.
- Qiusheng Huang, Xiaohui Zhong, Xu Fan, and Hao Li. FuXi-RTM: A physics-guided prediction framework with radiative transfer modeling. In *Proceedings of the IEEE/CVF International Conference on Computer Vision*, pp. 8852–8862, 2025.
- Remi Lam, Alvaro Sanchez-Gonzalez, Matthew Willson, et al. Learning skillful medium-range global weather forecasting. *Science*, 382(6677):1416–1421, 2023.
- Simon Lang, Mihai Alexe, Matthew Chantry, et al. Aifs – ecmwf's data-driven forecasting system. *arXiv preprint arXiv:2406.01465*, 2024.
- Tung Nguyen, Johannes Brandstetter, Archit Kapoor, Jayesh K. Gupta, and Aditya Grover. Climax: A foundation model for weather and climate. In *Proceedings of the 40th International Conference on Machine Learning (ICML)*, 2023.
- Jaideep Pathak, Shashank Subramanian, Peter Harrington, et al. Fourcastnet: A global data-driven high-resolution weather model using adaptive fourier neural operators. *arXiv preprint arXiv:2202.11214*, 2022.
- Ilan Price, Alvaro Sanchez-Gonzalez, Ferran Alet, et al. Probabilistic weather forecasting with machine learning. *Nature*, 637(8044):84–90, 2025. doi: 10.1038/s41586-024-08252-9.
- Stephane Vannitsem, John Bjornar Bremnes, Jonathan Demaeyer, et al. Statistical postprocessing for weather forecasts: Review, challenges, and avenues in a big data world. *Bulletin of the American Meteorological Society*, 102(3):E681–E699, 2021.
- Zesheng Ye, Chengyi Cai, Ruijiang Dong, Jianzhong Qi, Lei Feng, Pin-Yu Chen, and Feng Liu. Neural network reprogrammability: A unified theme on model reprogramming, prompt tuning, and prompt instruction, 2025. *arXiv:2506.04650*.
- Zeping Zhang, Wei Han, Lei Chen, Hao Li, Lipeng Jiang, and Bin Zhao. Fine-tuning fuxi with cma's reanalysis data to improve forecasting. *Journal of Meteorological Research*, 39(6):1411–1424, 2025. doi: 10.1007/s13351-025-5052-y.

Table 3: A summary of the 70 channels used by WeatherBridge. The Units column gives the physical unit of each channel.

Type	Full name	Abbreviation	Role	Units	Channels
Upper-air variables	Geopotential	Z	Input and output	$\text{m}^2 \text{s}^{-2}$	13
	Temperature	T	Input and output	K	13
	u component of wind	U	Input and output	m s^{-1}	13
	v component of wind	V	Input and output	m s^{-1}	13
	Relative humidity	R	Input and output	%	13
Surface variables	2-m temperature	T2M	Input and output	K	1
	10-m u wind component	U10M	Input and output	m s^{-1}	1
	10-m v wind component	V10M	Input and output	m s^{-1}	1
	Mean sea-level pressure	MSL	Input and output	Pa	1
	Total precipitation	TP	Input and output	m	1
Total					70

Table 4: Per-channel reference error $\text{RMSE}_{\text{ref}}^g$ used to normalize the ORA loss: the RMSE of the frozen backbone initialized from ERA5 at the first inference step.

Channel	Per-channel RMSE
Z500	21.8412
Z850	22.2562
U200	1.0605
T2M	0.6205
U10M	0.4522
V10M	0.4659
MSL	31.0368
TP	0.6198
T300	0.2347

A IMPLEMENTATION AND EVALUATION DETAILS

A.1 TRAINING VARIABLES LIST

Table 3 summarizes the 70 channels used by our model: 65 upper-air channels comprising five variables (z , t , u , v , r) at 13 pressure levels (50, 100, 150, 200, 250, 300, 400, 500, 600, 700, 850, 925, and 1000 hPa), and five surface variables (t_{2m} , u_{10m} , v_{10m} , msl , tp). All source and ERA5 fields are normalized using the same ERA5 per-channel means and standard deviations. The wind speeds ws_{10} and ws_{850} are computed as the magnitudes of the horizontal wind vectors from the corresponding U and V components at 10 m and 850 hPa, respectively.

A.2 CHANNEL LIST AND NORMALIZATION

Table 4 lists the nine guided channels of the ORA loss and the per-channel reference error $\text{RMSE}_{\text{ref}}^g$ used to normalize them; the normalization is defined in Section 3.3.

A.3 EVALUATION METRICS

$$\text{RMSE}_c(\tau) = \sqrt{\frac{\sum_i \cos(\phi_i) (f_{c,i}(\tau) - o_{c,i}(\tau))^2}{\sum_i \cos(\phi_i)}} \quad (9)$$

$$\text{ACC}_c(\tau) = \frac{\sum_i \cos(\phi_i) f'_{c,i}(\tau) o'_{c,i}(\tau)}{\sqrt{\sum_i \cos(\phi_i) f'^2_{c,i}(\tau) \cdot \sum_i \cos(\phi_i) o'^2_{c,i}(\tau)}} \quad (10)$$

Table 5: ERA5 reference initialization: the frozen backbone driven by ERA5 fields, per evaluation variable, at 6 h and 240 h.

Variable	RMSE (6 h)	ACC (6 h)	RMSE (240 h)	ACC (240 h)
Z500	21.71	0.9996	745.90	0.5420
T2M	0.60	0.9825	2.67	0.5937
T850	0.38	0.9943	3.49	0.4757
WS10	0.46	0.9841	2.97	0.2774
WS850	0.66	0.9892	5.10	0.2853
MSL	32.02	0.9990	686.49	0.4975

RMSE and ACC are computed per variable, per lead time and per source; $\cos(\phi)$ provides the latitude weighting. Here f and o are the forecast and ERA5 truth fields, f' and o' are anomaly fields relative to the climatology, τ is the lead time, and $\cos(\phi)$ provides area weighting.

Lower RMSE indicates smaller errors; higher ACC indicates better agreement in anomaly patterns. Following FuXi-RTM (Huang et al., 2025), we express the RMSE difference of WeatherBridge relative to the baseline as

$$\text{NRMSE}_c^s(n) = \frac{\text{RMSE}_c^s(n) - \text{RMSE}_{c,\text{base}}^s(n)}{\text{RMSE}_{c,\text{base}}^s(n)} \times 100\%. \quad (11)$$

Thus, negative NRMSE indicates improvement over the baseline.

PSNR and SSIM assess the aligned initial field $\hat{x}_t^E$ against the ERA5 field x_t^E at the same valid time, per variable and per source, and are computed over the whole field with a single window, without latitude weighting:

$$\text{PSNR}_c = 20 \log_{10} \left(\frac{L_c}{\sqrt{\frac{1}{N} \sum_i (\hat{x}_{c,i}^E - x_{c,i}^E)^2}} \right) \quad (12)$$

$$\text{SSIM}_c = \frac{(2\mu_{\hat{x}}\mu_x + C_1)(2\sigma_{\hat{x}x} + C_2)}{(\mu_{\hat{x}}^2 + \mu_x^2 + C_1)(\sigma_{\hat{x}}^2 + \sigma_x^2 + C_2)} \quad (13)$$

where μ and σ are the spatial mean and standard deviation of the aligned field $\hat{x}$ and of the ERA5 field x , $\sigma_{\hat{x}x}$ is their spatial covariance, and N is the number of grid points. L_c is the data range of variable c , defined as the difference between the 1st and 99th percentiles of the ERA5 field, and it also sets the SSIM constants $C_1 = (0.01 L_c)^2$ and $C_2 = (0.03 L_c)^2$.

A.4 ERA5 REFERENCE INITIALIZATION

Table 5 reports the reference path, in which the frozen backbone is driven by ERA5 fields, per evaluation variable. It is the best initialization available in this setup and therefore serves as the reference against which raw-source and WeatherBridge-aligned initialization are compared.

B EXTENDED RESULTS

B.1 STATISTICAL SIGNIFICANCE OF THE MAIN RESULTS

We also performed significance tests on the improvements reported in Section 4.2. For each of the 326 daily 00z initializations shared between the raw-source and WeatherBridge-aligned runs, we compute the mean over the 40 six-hourly lead steps, form the per-initialization relative improvement, and bootstrap its mean (10,000 resamples) to obtain a 95% percentile confidence interval (Table 6). The intervals exclude zero for 15 of the 18 source-variable pairs on RMSE and for 13 of the 18 pairs on ACC: all six GFS and all six CMA variables are significant, whereas among the HRES variables only T2M, T850 and WS10 reach significance on RMSE, consistent with HRES already starting close to the ERA5 reference and therefore having little headroom. The three RMSE pairs whose interval contains zero (Z500, WS850 and MSL on HRES) differ by less than 0.2% in the mean.

Table 6: Paired bootstrap confidence intervals for the mean rollout RMSE improvement over the raw-source initialization. Each initialization’s mean over the 40 six-hourly lead steps (6 h to 240 h) is formed first and the improvement is then averaged over the 326 daily 00z initializations; the last column marks whether the 95% percentile interval (10,000 resamples) excludes zero. The pooled mean in Table 1 weights every lead step equally instead, so the two averages differ slightly. ACC behaves the same way: 13 of the 18 pairs have an interval excluding zero.

	Source	Var	Mean RMSE improvement (%)	95% CI	Sig.
GFS		Z500	+46.11	[+45.50, +46.75]	yes
		T2M	+45.70	[+45.14, +46.24]	yes
		T850	+41.21	[+40.76, +41.65]	yes
		WS10	+32.60	[+32.30, +32.91]	yes
		WS850	+30.54	[+30.20, +30.88]	yes
		MSL	+41.75	[+41.12, +42.40]	yes
HRES		Z500	-0.17	[-0.64, +0.28]	no
		T2M	+0.72	[+0.50, +0.95]	yes
		T850	+0.57	[+0.33, +0.80]	yes
		WS10	+0.17	[+0.00, +0.33]	yes
		WS850	-0.00	[-0.18, +0.17]	no
		MSL	+0.18	[-0.26, +0.61]	no
CMA		Z500	+9.17	[+8.15, +10.18]	yes
		T2M	+25.87	[+25.11, +26.61]	yes
		T850	+9.97	[+9.39, +10.54]	yes
		WS10	+5.72	[+5.37, +6.06]	yes
		WS850	+4.14	[+3.80, +4.48]	yes
		MSL	+19.35	[+18.29, +20.41]	yes

Table 7: Ablation study. Each variant is compared with the full model trained on the same sources: the ORA-weight ablation is trained on GFS only, and the source-embedding ablation is trained on GFS, HRES and CMA.

Source	Setting	SSIM	PSNR	RMSE (6 h) ↓						ORA loss
				Z500	T2M	T850	WS10	WS850	MSL	
ORA weight ablation (GFS only)										
GFS	$\lambda_f = 8 \times 10^{-3}$ (default)	0.9839	32.70	42.20	1.04	0.91	0.88	1.51	70.98	1.8491
	$\lambda_f = 0$	0.9844	32.57	45.43	1.08	0.90	0.90	1.49	81.67	1.9702
Source-embedding ablation (GFS, HRES and CMA)										
GFS	Full WeatherBridge	0.9851	32.99	42.07	1.01	0.88	0.88	1.49	69.20	1.7574
	w/o source embedding	0.9851	32.97	40.20	1.03	0.87	0.91	1.46	74.73	1.7718
HRES	Full WeatherBridge	0.9904	35.81	29.39	0.81	0.70	0.72	1.19	46.80	1.7574
	w/o source embedding	0.9903	35.76	29.33	0.82	0.71	0.72	1.20	47.58	1.7718
CMA	Full WeatherBridge	0.9827	32.18	53.85	1.10	1.03	0.93	1.63	77.26	1.7574
	w/o source embedding	0.9826	32.12	53.49	1.11	1.04	0.93	1.63	77.78	1.7718

B.2 ABLATION STUDY

In this experiment setting the ORA weight to zero is equivalent to w/o ORA loss, and it degrades the rollout: with both models trained on GFS only, the 6 h RMSE increases for 4 of the 6 evaluation variables, by 2.0% to 15.1%, and changes by less than 1% for the remaining two (T850 and WS850, where it slightly decreases), and the measured ORA loss rises from 1.8491 to 1.9702. Removing the source embedding also degrades the rollout: the 6 h RMSE increases for 13 of the 18 source-variable pairs, by up to 8.0%, and its contribution is largest under source imbalance, where removing it raises the 6 h RMSE for 17 of the 18 pairs and by up to 13.9% (Tables 7 and 8).

Degradation under imbalance scales with the training share: for the scarce CMA source, PSNR/SSIM drop from 32.18/0.9827 (balanced) to 31.73/0.9808 (imbalanced) and further to 31.14/0.9782 when the source embedding is removed (imb_wo_src), while the dominant HRES source is least affected (PSNR 35.81, 35.71, 35.47). Removing the embedding under imbalance degrades 17 of the 18 source-variable pairs at 6 h, by up to 13.9%, so the source embedding con-

Table 8: Source-imbalance stress test: balanced full-source training, imbalanced training (HRES 100%, GFS 50%, CMA 10%), and imbalanced training without the source embedding. PSNR/SSIM describe the WeatherBridge-aligned initial field against ERA5, so they do not depend on lead time and are listed once, in the upper block; the upper block then gives the RMSE of the six evaluation variables at the first step (6 h) and the lower block the same RMSE at 240 h, where forecast error has largely saturated.

Source	Training mix	PSNR	SSIM	RMSE (6 h) ↓					
				Z500	T2M	T850	WS10	WS850	MSL
GFS	Balanced	32.99	0.9851	42.07	1.01	0.88	0.88	1.49	69.20
	Imbalanced	32.62	0.9840	42.04	1.03	0.91	0.89	1.50	71.13
	Imbalanced, w/o src emb	31.94	0.9823	43.55	1.12	0.97	0.91	1.53	79.29
HRES	Balanced	35.81	0.9904	29.39	0.81	0.70	0.72	1.19	46.80
	Imbalanced	35.71	0.9900	28.89	0.82	0.71	0.72	1.20	46.97
	Imbalanced, w/o src emb	35.47	0.9897	29.90	0.86	0.76	0.74	1.22	49.96
CMA	Balanced	32.18	0.9827	53.85	1.10	1.03	0.93	1.63	77.26
	Imbalanced	31.73	0.9808	53.24	1.17	1.06	0.95	1.67	79.17
	Imbalanced, w/o src emb	31.14	0.9782	51.19	1.33	1.15	0.99	1.70	84.00
RMSE (240 h) ↓									
GFS	Balanced			790.22	2.78	3.65	3.05	5.23	727.50
	Imbalanced			788.96	2.78	3.64	3.05	5.24	727.58
	Imbalanced, w/o src emb			784.68	2.80	3.63	3.04	5.21	722.84
HRES	Balanced			747.00	2.68	3.48	2.98	5.10	691.29
	Imbalanced			746.09	2.68	3.47	2.98	5.11	691.46
	Imbalanced, w/o src emb			747.26	2.68	3.48	2.98	5.11	692.05
CMA	Balanced			806.73	2.83	3.71	3.07	5.28	742.12
	Imbalanced			798.97	2.82	3.69	3.07	5.28	736.94
	Imbalanced, w/o src emb			802.00	2.84	3.70	3.08	5.29	741.08

tributes most when the sources are imbalanced (Table 8). By 240 h these differences have largely disappeared: the three training settings differ by at most 1.0% on any source-variable pair, and the ordering seen at 6 h no longer holds, so the imbalance penalty is an early-rollout effect that error growth masks at long lead times. Adding the source embedding leaves 5 of the 18 source-variable pairs slightly worse, but it lowers the overall ORA loss, which averages the per-channel RMSE normalized by the ERA5 reference error over the guided channels, from 1.7718 to 1.7574. The embedding therefore favours the aggregate multi-variable error over any single variable, so individual variables may move in either direction while the overall error still decreases.

B.3 PARAMETER SENSITIVITY

Increasing the ORA weight from 0 to the default $8e-3$ lowers the first-step Z500 RMSE from 45.43 to 42.20 while Z500 ACC stays around 0.999; SSIM/PSNR peak near $2e-3$ (0.9845, 32.81) and degrade sharply beyond $1.28e-1$ (SSIM 0.9790, PSNR 30.75). The default $8e-3$ therefore sits on the fidelity-utility frontier (Figure 6).

B.4 MULTI-SOURCE TRAINING STRUCTURE (SOURCE MIXING)

Different mixtures of sources give different results (Table 9). Evaluated on a given source, training together with a second source usually beats training on that source alone: for GFS, adding HRES cuts the 6 h RMSE of the most improved variable, T850, by 2.2% and degrades none of the six, for a mean reduction of 1.4%, while adding CMA instead leaves three of the six variables slightly worse, by at most 1.1%, and the mean 0.4% higher. Over all sources the worst pair is still only 2.9% behind the single-source model. The gain is largest for CMA, the weakest source on its own, which improves on five of its six variables when trained together with HRES. We therefore train the final adapter on all three sources, since no negative interference between sources is observed.




Figure 6: Sensitivity to the ORA weight λ_f (default 8×10^{-3} ; source: GFS). Each panel shows one metric against the weight: initial-field SSIM and PSNR, first-step Z500 RMSE and ACC, and the ORA loss; the star marks the default setting.

Table 9: Multi-source training structure: single-source models and pair-trained adapters, each evaluated on every source.

Eval source	Training setup	PSNR	SSIM	RMSE (6 h) $\downarrow$					
				Z500	T2M	T850	WS10	WS850	MSL
GFS	GFS-only	32.70	0.9839	42.20	1.04	0.91	0.88	1.51	70.98
	GFS+CMA	32.67	0.9842	42.51	1.04	0.91	0.89	1.51	71.32
	GFS+HRES	32.82	0.9844	41.35	1.03	0.89	0.88	1.49	69.60
HRES	HRES-only	35.67	0.9898	29.01	0.82	0.72	0.72	1.21	47.06
	GFS+HRES	35.72	0.9901	29.28	0.82	0.72	0.72	1.20	47.17
	CMA+HRES	35.81	0.9904	29.16	0.81	0.71	0.72	1.19	46.72
CMA	CMA-only	31.87	0.9812	51.99	1.17	1.06	0.95	1.66	77.62
	GFS+CMA	31.95	0.9815	53.49	1.14	1.05	0.94	1.66	78.03
	CMA+HRES	32.17	0.9828	53.06	1.10	1.03	0.93	1.62	77.07

B.5 DATA SCALE

Rollout and initial-field metrics improve monotonically with more paired data: SSIM 0.9795 to 0.9839, PSNR 31.12 to 32.70, and the 6 h RMSE of every evaluation variable falls from 25% to 100% of the samples, for example from 44.45 to 42.20 for Z500 and from 83.97 to 70.98 for MSL, showing that the source-to-ERA5 structured bias is learnable and that WeatherBridge scales with data (Table 10).

B.6 EFFICIENCY AND TRAINING COST

The smallest configuration is the best: Lite (138.4M) attains the lowest 6 h RMSE on all six evaluation variables and the lowest ORA loss (1.8491), while Mid (222.5M) and Deep (285.4M) are slightly worse (for example from 1.04 to 1.08 and 1.11 for T2M, and from 70.98 to 71.89 and 74.35 for MSL). Capacity scaling therefore saturates, which supports using the Lite model as the main configuration and treating WeatherBridge as a lightweight front-end adapter (Table 11). The Lite model trains in about 88 h on 4 A100 GPUs (about 350 GPU-hours), and the test-time FuXi rollout over the three sources (326 initializations each) takes about 46 h in total.

Table 10: Data-scale study (default source GFS): training on 25%, 50% and 100% of the paired samples.

Training samples	SSIM	PSNR	RMSE (6 h) ↓						ORA loss
			Z500	T2M	T850	WS10	WS850	MSL	
25%	0.9795	31.12	44.45	1.20	1.03	0.92	1.54	83.97	2.0340
50%	0.9803	31.45	44.46	1.17	1.00	0.92	1.54	83.57	2.0160
100%	0.9839	32.70	42.20	1.04	0.91	0.88	1.51	70.98	1.8491

Table 11: Model-scale study (default source: GFS).

Model	SSIM	PSNR	RMSE (6 h) ↓						ORA loss
			Z500	T2M	T850	WS10	WS850	MSL	
Lite	0.9839	32.70	42.20	1.04	0.91	0.88	1.51	70.98	1.8491
Mid	0.9836	32.50	42.65	1.08	0.92	0.89	1.52	71.89	1.8793
Deep	0.9831	32.33	43.25	1.11	0.94	0.90	1.52	74.35	1.8986

Table 12: Naive and WeatherBridge-aligned initialization on FengWu for the GFS source: RMSE and ACC at 6 h and 240 h per variable, with the raw-source value followed by the WeatherBridge-aligned value and by Δ , the relative improvement of the WeatherBridge-aligned result over the raw-source result in percent, computed before rounding; “↑” and “↓” indicate that higher and lower values are better, respectively.

Var	RMSE 6 h ↓			ACC 6 h ↑			RMSE 240 h ↓			ACC 240 h ↑		
	raw	WeatherBridge	Δ	raw	WeatherBridge	Δ	raw	WeatherBridge	Δ	raw	WeatherBridge	Δ
Z500	226.94	56.41	75.14%	0.9621	0.9976	3.70%	1023.32	952.65	6.91%	0.0732	0.1748	138.68%
T2M	2.28	1.22	46.24%	0.7485	0.9268	23.82%	3.93	3.23	17.77%	0.3584	0.4365	21.78%
T850	6.14	0.93	84.83%	0.2846	0.9645	238.90%	4.81	4.08	15.25%	0.0783	0.1717	119.27%
WS10	1.65	0.90	45.49%	0.7880	0.9381	19.05%	3.88	3.86	0.61%	0.0479	0.0907	89.50%
WS850	2.77	1.56	43.57%	0.7948	0.9383	18.06%	6.29	5.91	6.08%	0.0322	0.0873	171.17%
MSL	297.15	80.75	72.82%	0.9112	0.9938	9.07%	867.90	727.29	16.20%	0.0952	0.2912	205.81%

B.7 TRANSFER TO A SECOND BACKBONE (FENGWU)

WeatherBridge is trained jointly with a frozen FuXi backbone. In this experiment the FengWu backbone is used as released: only its initial fields are replaced by the WeatherBridge-aligned states, so the adapter is applied unchanged and nothing is retrained for FengWu. Table 2 reports the resulting rollout on the GFS source, which shows that the adapted states are usable by a different ERA5-trained backbone.